

PE/Cyanine5 anti-human CD161

Catalog # / Size: 2299750 / 25 tests
2299755 / 100 tests

Clone: HP-3G10

Isotype: Mouse IgG1, κ


Immunogen: Human NK cells

Reactivity: Human, Non-human primate, Other

Preparation: The antibody was purified by affinity chromatography and conjugated with PE/Cyanine5 under optimal conditions.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA (origin USA)

Concentration: Lot-specific



Human peripheral blood lymphocytes were stained with FITC anti-human CD4 and PE/Cyanine5 anti-human CD161 (clone HP-3G10) (left) or PE/Cyanine5 mouse IgG1, κ isotype control (right).

Applications:

Applications: Flow Cytometry

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is 5 μ L per million cells in 100 μ L staining volume or 5 μ L per 100 μ L of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

Application Notes: Additional reported applications (for the relevant formats) include: inhibition of cytokine production and Western blotting under nonreducing conditions.

- Application References:**
1. Gumß M, et al. 2004. *Blood* 104:3664.
 2. Exley M, et al. 1998. *J. Exp. Med.* 188:867.
 3. Marquez C, et al. 1998. *Blood* 91:2760.

Description: CD161 is a type II transmembrane glycoprotein, also known as NKR-P1A, that is expressed as a 40-44 kD homodimer. It is a member of the C-type lectin superfamily. CD161 is expressed on a majority of NK cells, NKT cells, and subsets of peripheral T cells and CD3⁺ thymocytes. It has been reported that Th17 cells are a subpopulation of CD4⁺CD161⁺CCR6⁺ cells. While the biological function of CD161 is not clear, it has been suggested to serve either as a stimulatory receptor or to inhibit NK cell-mediated cytotoxicity and cytokine production. LLT-1 (lectin-like transcript-1, also named as osteoclast inhibitory lectin or CLEC2D) is the ligand of CD161.

- Antigen References:**
1. Takahashi T, et al. 2006. *J. Immunol.* 176:211.
 2. Cosmi L, et al. 2008. *J. Exp. Med.* 205:1903.
 3. Aldemir H, et al. 2005. *J. Immunol.* 175:7791.
 4. Rosen DB, et al. 2008. *J. Immunol.* 180:6508.