SONY

Product Data Sheet

APC anti-human HLA-E

Catalog # / Size: 2313030 / 100 tests
2313025 / 25 tests

Clone: 3D12

Isotype: Mouse IgG1, κ

Reactivity: Human

Preparation: The antibody was purified by affinity chromatography and conjugated with APC under optimal conditions.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and BSA (origin USA).

Workshop Number: HCDM listed

Concentration: Lot-specific



Human peripheral blood lymphocytes were stained with HLA-E (clone 3D12) APC (filled histogram) or mouse IgG1, κAPC isotype control (open histogram).

Applications:

Applications: Flow Cytometry

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is 5 µl per million cells or 5 µl per 100 µl of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

Human paraffin-embedded colon tissue slices were prepared with a standard protocol of deparaffinization and rehydration. Antigen retrieval was done with Citrate buffer 1X pH 6.0 at 95°C for 40 minutes. Tissue was washed with PBS/0.05% Tween 20 twice for five minutes and blocked with 5% FBS and 0.2% gelatin for 30 minutes. Then, the tissue was stained with 10 µg/mL of anti-human CD66a/c/e (clone ASL-32) Spark YG™ 570 (red) and anti-Pan-Cytokeratin (clone C-11) Alexa Fluor® 488 antibody (green) at 4°C overnight. Nuclei were counterstained with DAPI (blue). The image was scanned with a 10X objective and stitched with software.

For research use only. Not for diagnostic use. Not for resale. Sony Biotechnology Inc. will not be held responsible for patent infringement or other violations that may occur with the use of our products.
Sony Biotechnology Inc. 1730 North First Street, San Jose, CA 95112 www.sonybiotechnology.com

Application References:

1. Lee N, *et al.* 1998. *Proc. Natl. Acad. Sci. USA.* 95:5199.
2. Wooden SL, *et al.* 2005. *J. Immunol.* 175:1383.
3. Monaco EL, *et al.* 2008. *J. Immunol.* 181:5442.
4. Corrah TW, *et al.* 2011. *J. Virol.* 85:3367. PubMed

Description: HLA-E is a non-classical MHC class I (MHC-Ib) molecule. It is characterized by limited polymorphism and ubiquitous expression. HLA-E, as other MHC-I molecules, is heterodimerized with β2-microglobulin. HLA-E interacts with CD94/NKG2 receptor to regulate NK cell cytolysis activities.

Antigen References:

1. Sullivan LC, *et al.* 2008. *Tissue Antigen.* 72:415.
2. Koller BH, *et al.* 1988. *J. Immunol.* 141:897.

For research use only. Not for diagnostic use. Not for resale. Sony Biotechnology Inc. will not be held responsible for patent infringement or other violations that may occur with the use of our products.