SONY

Product Data Sheet

APC/Fire™ 750 anti-human TCR Vα24-Jα18 (iNKT cell)

Catalog # / Size: 2314635 / 25 tests
2314640 / 100 tests

Clone: 6B11

Isotype: Mouse IgG1, κ

Reactivity: Human, Other

Preparation: The antibody was purified by affinity chromatography and conjugated with APC/Fire™ 750 under optimal conditions.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA (origin USA).

Concentration: Lot-specific



Human peripheral blood lymphocytes were stained with CD3 FITC and anti-human TCR V7α24-Jα18 (clone 6B11) APC/Fire™ 750 (left) or mouse IgG1, κ APC/Fire™ 750 isotype control (right).

Applications:

Applications: Flow Cytometry

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is 5 µL per million cells in 100 µL staining volume or 5 µL per 100 µL of whole blood.

* APC/Fire™ 750 has a maximum excitation of 650 nm and a maximum emission of 787 nm.

Application Notes: The 6B11 antibody recognizes the invariant CDR3 region of TCR Va24-JaQ.

Application References:

1. Rout N, *et al.* 2010. *PLoS One* 5:e9787. (FC)

Description: Encoded by the TCR Vα24-Jα18 germline configuration, Vα24-JαQ is expressed on a subset of NKT cells, namely invariant NKT (iNKT). Vα24-JαQ TCR interacts with the glycolipid loaded MHC class 1b molecule CD1d, inducing activation and subsequent cytokine production. iNKT cells have been implicated in immune regulation, tumor surveillance, and host response to pathogens. While iNKT cells occur at low frequency in the blood, assorted chemokines contribute to their tissue homing potential.

Antigen References:

1. Thomas SY, *et al.* 2003. *J. Immunol.* 171:2571.
2. Exley MA, *et al.* 2008. *Eur. J. Immunol.* 38:1756.
3. Montoya CJ, *et al.* 2007. *Immunology.* 122:1.
4. Gansuvd B, *et al.* 2003. *J. Immunol.* 171:2904.