


PE anti-human CD209 (DC-SIGN)

Catalog # / Size: 2315020 / 100 tests
Clone: DCS-8C1
Isotype: Mouse IgG2b, κ
Immunogen: Extracellular domain of human DC-SIGN
Reactivity: Human
Preparation: The antibody was purified by affinity chromatography, and conjugated with PE under optimal conditions. The solution is free of unconjugated PE and unconjugated antibody.
Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA (origin USA).
Concentration: Lot-specific



Human monocyte-derived dendritic cells stained with DCS-8C1 PE

Applications:

Applications: Flow Cytometry
Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. **Test size products are transitioning from 20 microL to 5 microL per test.** Please check your vial or your CoA to find the suggested use of this reagent per million cells in 100 microL staining volume or per 100 microL of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

Description: CD209, known as Dendritic Cell-Specific Intercellular adhesion molecule 3 (ICAM-3)-Grabbing Nonintegrin (DC-SIGN), is a 44 kD type II transmembrane glycoprotein and a member of the C-type lectin family. CD209 is expressed on myeloid dendritic cells, placental macrophages, liver and placental endothelial cells. CD209 binds to ICAM-3 (CD50), ICAM-2 (CD102), and Butyrophilin (BTN2A1), and mediates dendritic cell migration and T cell proliferation. Importantly, CD209 is a receptor of HIV-1 and some other viruses (such as West Nile virus, hepatitis C virus, etc), and some bacteria or parasites. It plays a critical role in capturing and internalizing those pathogens. LSP1 (leukocyte-specific protein 1) interacts with the cytoplasmic domain of CD209 and mediates transport of HIV to the proteasome.

Antigen References: 1. Granelli-Piperno A, *et al.* 2005. *J Immunol.* 175:4265.