

Pacific Blue™ anti-human CD34

Catalog # / Size: 2317555 / 25 µg
2317560 / 100 µg

Clone: 581

Isotype: Mouse IgG1, κ


Reactivity: Human

Preparation: The antibody was purified by affinity chromatography, and conjugated with Pacific Blue™ under optimal conditions. The solution is free of unconjugated Pacific Blue™.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.

Workshop Number: V MA27

Concentration: 0.5




Human peripheral blood leukocytes stained with CD34 Pacific Blue™ (clone 581) or mouse IgG1, κ Pacific Blue™ isotype control and costained with CD45 FITC, CD14 PE and 7-AAD, then analyzed via gating on total live and CD14 negative lymph

Applications:

Applications: Flow Cytometry

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is ≤ 1.0 microg per 10⁶ cells in 100 microL volume or 100 microL of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.



* Pacific Blue™ has a maximum emission of 455 nm when it is excited at 405 nm. Prior to using Pacific Blue™ conjugate for flow cytometric analysis, please verify your flow cytometer's capability of exciting and detecting the fluorochrome.

Application Notes: The 581 antibody recognizes the class III group epitope which is resistant to sialidase/glycolyprotease and chymopapain treatment. Additional reported applications (for the relevant formats) include: immunohistochemical staining of paraffin-embedded tissue sections⁵ and immunofluorescence⁶.

Application References: 1. Schlossman SF, et al. 1995. *Leukocyte Typing V: White Cell Differentiation Antigen*. New York: Oxford University Press.

2. Felschow DM, *et al.* 2001. *Blood* 97:3768.
 3. Rudin CE, *et al.* 1997. *Br. J. Haematol.* 97:488.
 4. Yoshino N, *et al.* 2000. *Exp. Anim. (Tokyo)* 49:97. (FC)
 5. Skowasch D, *et al.* 2003. *Cardiovasc Res.* 60:684. (IHC)
 6. Umland O, *et al.* 2003. *J. Histochem. Cytochem.* 51:977. (IF)
 7. Bhatwadekar AD, *et al.* 2010. *Diabetes.* 59:2010. [PubMed](#)
 8. Jarajapu YP, *et al.* 2013. *Diabetes.* 62:1258. [PubMed](#).
-

Description: CD34, also known as gp105-120, is a type I monomeric sialomucin-like glycoprophosphoprotein with an approximate molecular weight of 105-120 kD. Selectively expressed on the majority of hematopoietic stem/progenitor cells, bone marrow stromal cells, capillary endothelial cells, embryonic fibroblasts, and some nervous tissue, CD34 is a commonly used marker to identify human hematopoietic stem/progenitor cells. According to the differential sensitivity to enzymatic cleavage, four groups of epitopes of CD34 have been described. CD34 mediates cell adhesion and lymphocytes homing through binding to L-selectin and E-selectin ligands.

Antigen
References:

1. Krause DS, *et al.* 1996. *Blood* 87:1.
2. Puri KD, *et al.* 1995. *J. Cell Biol.* 131:261.
3. Zola H, *et al.* 2007. *Leukocyte and Stromal Cell Molecules: The CD Markers.* John Wiley & Sons In