

Pacific Blue™ anti-human CD73 (Ecto-55'-nucleotidase)

Catalog # / Size: 2320055 / 25 tests
2320060 / 100 tests

Clone: AD2

Isotype: Mouse IgG1, κ


Reactivity: Human

Preparation: The antibody was purified by affinity chromatography and conjugated with Pacific Blue™ under optimal conditions. The solution is free of unconjugated Pacific Blue™.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA (origin USA).

Workshop Number: V B-CD73.3

Concentration: Lot-specific




Human peripheral blood lymphocytes were stained with CD3 FITC and CD73 (clone AD2) Pacific Blue™ (top) or mouse IgG1, κ Pacific Blue™ isotype control (bottom).

Applications:

Applications: Flow Cytometry

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is 5 microL per million cells or 5 microL per 100 microL of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.



* Pacific Blue™ has a maximum emission of 455 nm when it is excited at 405 nm. Prior to using Pacific Blue™ conjugate for flow cytometric analysis, please verify your flow cytometer's capability of exciting and detecting the fluorochrome.

Application Notes: Additional reported applications (for the relevant formats) include: immunofluorescence3.

Application References: 1. Nakamura T, *et al.* 1993. *J. Immunol.* 151:6933.
2. Liao J, *et al.* 2011. *J Endod.* 37:1217. [PubMed](#)
3. Touboul C, *et al.* 2013. *J. Transl. Med.* 11:28. (IF)

Description: CD73 is a 70 kD glycoposphatidylinositol (GPI)-linked 5'-nucleotidase, which is also known as ecto-5'-nucleotidase. It converts adenosine monophosphate (AMP) to adenosine. CD73 is expressed on subsets of T and B cells, mesenchymal stem

cells, follicular dendritic cells, endothelial cells, and epithelial cells. It has been reported that CD73 costimulates T cell activation, and mediates adhesion of lymphocytes to follicular dendritic cells and endothelial cells.

**Antigen
References:**

1. Zola H, *et al.* 2007. *Leukocyte and stromal Cell Molecules:the CD Markers.* A John Wiley & Sons Inc, Publication.
2. Airas L and Jalkanen S, *et al.* 1996. *Blood* 88:1755.
3. Gutensohn W, *et al.*