

APC/Cy7 anti-human CD73 (Ecto-55'-nucleotidase)

Catalog # / Size: 2320105 / 25 tests
2320110 / 100 tests

Clone: AD2

Isotype: Mouse IgG1, κ


Reactivity: Human

Preparation: The antibody was purified by affinity chromatography and conjugated with APC/Cy7 under optimal conditions. The solution is free of unconjugated APC/Cy7 and unconjugated antibody.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA (origin USA).

Workshop Number: V B-CD73.3


Concentration: Lot-specific



Applications:

Applications: Flow Cytometry

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is 5 microL per million cells or 5 microL per 100 microL of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.



This product is subject to proprietary rights of Sirigen Inc. and is made and sold under license from Sirigen Inc. The purchase of this product conveys to the buyer a non-transferable right to use the purchased product for research purposes only. This product may not be resold or incorporated in any manner into another product for resale. Any use for therapeutics or diagnostics is strictly prohibited. This product is covered by U.S. Patent(s), pending patent applications and foreign equivalents.

Human peripheral blood lymphocytes were stained with CD3 FITC and CD73 (clone AD2) APC/Cy7 (top) or mouse IgG1, κ APC/Cy7 isotype control (bottom).

Application Notes: Additional reported applications (for the relevant formats) include:immunofluorescence3.

Application References: 1. Nakamura T, *et al.* 1993. *J. Immunol.* 151:6933.
2. Liao J, *et al.* 2011. *J Endod.* 37:1217. [PubMed](#)
3. Touboul C, *et al.* 2013. *J. Transl. Med.* 11:28. (IF)

Description: CD73 is a 70 kD glycoposphatidylinositol (GPI)-linked 5'-nucleotidase, which is also known as ecto-5'-nucleotidase. It converts adenosine monophosphate (AMP) to adenosine. CD73 is expressed on subsets of T and B cells, mesenchymal stem cells, follicular dendritic cells, endothelial cells, and epithelial cells. It has been reported that CD73 costimulates T cell activation, and mediates adhesion of lymphocytes to follicular dendritic cells and endothelial cells.

Antigen
References:

1. Zola H, *et al.* 2007. *Leukocyte and stromal Cell Molecules:the CD Markers*. A John Wiley & Sons Inc, Publication.
2. Airas L and Jalkanen S, *et al.* 1996. *Blood* 88:1755.
3. Gutensohn W, *et al.*