

Brilliant Violet 785® anti-human CD73 (Ecto-5'-nucleotidase)

Catalog # / Size: 2320135 / 25 tests
2320140 / 100 tests

Clone: AD2

Isotype: Mouse IgG1, κ


Reactivity: Human

Preparation: The antibody was purified by affinity chromatography and conjugated with Brilliant Violet 785™ under optimal conditions. The solution is free of unconjugated Brilliant Violet 785™ and unconjugated antibody.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and BSA (origin USA).

Workshop Number: V B-CD73.3

Concentration: 0.5




Human peripheral blood lymphocytes were stained with CD3 PE and CD73 (clone AD2) Brilliant Violet 785™ (top) or mouse IgG1, κ Brilliant Violet 785™ isotype control (bottom).

Applications:

Applications: Flow Cytometry

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is ≤5 microL per million cells or 5 microL per 100 microL of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.



Brilliant Violet 785™ excites at 405 nm and emits at 785 nm. The bandpass filter 780/60 nm is recommended for detection, although filter optimization may be required depending on other fluorophores used. Be sure to verify that your cytometer configuration and software setup are appropriate for detecting this channel. Refer to your instrument manual or manufacturer for support. Brilliant Violet 785™ is a trademark of Sirigen Group Ltd.

Application Notes: Additional reported applications (for the relevant formats) include: immunofluorescence3.

Application References: 1. Nakamura T, *et al.* 1993. *J. Immunol.* 151:6933.
2. Liao J, *et al.* 2011. *J Endod.* 37:1217. [PubMed](#)
3. Touboul C, *et al.* 2013. *J. Transl. Med.* 11:28. (IF)

Description: CD73 is a 70 kD glycoposphatidylinositol (GPI)-linked 5'-nucleotidase, which is also known as ecto-5'-nucleotidase. It converts adenosine monophosphate (AMP) to adenosine. CD73 is expressed on subsets of T and B cells, mesenchymal stem cells, follicular dendritic cells, endothelial cells, and epithelial cells. It has been reported that CD73 costimulates T cell activation, and mediates adhesion of lymphocytes to follicular dendritic cells and endothelial cells.

Antigen 1. Zola H, *et al.* 2007. *Leukocyte and stromal Cell Molecules:the CD Markers*. A
References: John Wiley & Sons Inc, Publication.
2. Airas L and Jalkanen S, *et al.* 1996. *Blood* 88:1755.
3. Gutensohn W, *et al.*