

**Brilliant Violet 421™ anti-human CD4**

**Catalog # / Size:** 2323160 / 100 tests  
2323155 / 25 tests

**Clone:** SK3


**Isotype:** Mouse IgG1,  $\kappa$

**Reactivity:** Human

**Preparation:** The antibody was purified by affinity chromatography and conjugated with Brilliant Violet 421™ under optimal conditions. The solution is free of unconjugated Brilliant Violet 421™ and unconjugated antibody.

**Formulation:** Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and BSA (origin USA).

**Concentration:** Lot-specific



Human peripheral blood lymphocytes were stained with CD4 (clone SK3) Brilliant Violet 421™ (filled histogram) or mouse IgG1,  $\kappa$  Brilliant Violet 421™ isotype control (open histogram).

**Applications:**

**Applications:** Flow Cytometry

**Recommended Usage:** Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is  $\leq 5$  microL per million cells or 5 microL per 100 microL of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

Brilliant Violet 421™ excites at 405 nm and emits at 421 nm. The standard bandpass filter 450/50 nm is recommended for detection. Brilliant Violet 421™ is a trademark of Sirigen Group Ltd.

This product is subject to proprietary rights of Sirigen Inc. and is made and sold under license from Sirigen Inc. The purchase of this product conveys to the buyer a non-transferable right to use the purchased product for research purposes only. This product may not be resold or incorporated in any manner into another product for resale. Any use for therapeutics or diagnostics is strictly prohibited. This product is covered by U.S. Patent(s), pending patent applications and foreign equivalents.

**Application References:**

1. Evans RL, *et al.* 1981. *Immunol.* 78:544
2. Arno A *et al.* 1999. *J. Infect. Dis.* 180:56
3. Muech M, *et al.* 1997. *Blood* 89:1364
4. Wang L, *et al.* 2012. *Cytometry A.* 81:567. [PubMed](#)

**Description:** CD4, also known as T4, is a 55 kD single-chain type I transmembrane glycoprotein expressed on most thymocytes, a subset of T cells, and monocytes/macrophages. CD4, a member of the Ig superfamily, recognizes antigens associated with MHC class II molecules and participates in cell-cell interactions, thymic differentiation, and signal transduction. CD4 acts as a primary receptor for HIV, binding to HIV gp120. CD4 has also been shown to interact with IL-16.

- Antigen** 1. Center D *et al.* 1996. *Immunol. Today* 17:476.
- References:** 2. Gaubin M *et al.* 1996. *Eur. J. Clin. Chem. Clin. Biochem.* 34:723.