

PerCP/Cy5.5 anti-human CD8

Catalog # / Size: 2323545 / 25 tests
2323550 / 100 tests

Clone: SK1


Isotype: Mouse IgG1, κ

Reactivity: Human

Preparation: The antibody was purified by affinity chromatography, and conjugated with PerCP/Cy5.5 under optimal conditions. The solution is free of unconjugated PerCP/Cy5.5 and unconjugated antibody.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA (origin USA).

Concentration: Lot-specific



Human peripheral blood lymphocytes stained with SK1 PerCp/Cy5.5

Applications:

Applications: Flow Cytometry

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is 5 microL per million cells or 5 microL per 100 microL of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

Application Notes: Clone SK1 recognizes the α chain of CD8. Additional reported applications (for the relevant formats) include: immunohistochemistry of acetone-fixed frozen tissue sections and formalin-fixed paraffin-embedded sections^{6,7}. This clone was tested in-house and does not demonstrate utility for formalin-fixed paraffin-embedded (FFPE) human tonsil sections. However, there are references cited that indicate that this clone has been used successfully in other FFPE applications^{6,7}.

- Application References:**
1. Ledbetter JA, *et al.* 1981. *J. Exp. Med.* 153:310.
 2. Campanelli R, *et al.* 2002. *Intl. Immunol.* 14:39.
 3. Evans RL, *et al.* 1981. *Immunol.* 78:544.
 4. Wooldridge L, *et al.* 2005. *J. Bio. Chem.* 280:27491.
 5. Ch'el IL, *et al.* 2011. *J Exp Med.* 208:633. [PubMed](#)
 6. Carbone A, *et al.* 1999. *Blood* 93:2319. (IHC)
 7. Ahmed A, *et al.* 2001. *J. Pathol.* 193:383. (IHC)

Description: CD8a is a 32-34 kD type I glycoprotein. It forms a homodimer (CD8a/a) or heterodimer (CD8a/b) with CD8b. CD8, also known as T8 and Leu2, is a member of the immunoglobulin superfamily found on the majority of thymocytes, a subset of peripheral blood T cells, and NK cells (which express almost exclusively CD8a homodimers). CD8 acts as a co-receptor with MHC class I-restricted T cell receptors in antigen recognition and T cell activation and has been shown to play a role in thymic differentiation. Two domains in CD8a are important for function: the extracellular IgSF domain binds the α_3 domain of MHC class I and the cytoplasmic CXCP motif binds the tyrosine kinase p56 Lck.

Antigen References:

1. Barclay N, *et al.* 1993. *The Leucocyte Antigen FactsBook*. Academic Press Inc. San Diego.

