
## **Product Data Sheet**

## **Purified anti-human CD3**

Catalog # / Size:	2324005 / 25 μg 2324010 / 100 μg
Clone:	SK7
Isotype:	Mouse IgG1, к
<b>Reactivity:</b>	Human
Preparation:	The antibody was purified by affinity chromatography.
Formulation:	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.
<b>Concentration:</b>	0.5



Human peripheral blood lymphocytes stained with SK7 FITC.

## **Applications:**

Applications:	Immunofluorescence
Recommended Usage:	Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is $\leq 0.25$ microg per million cells in 100 microL volume. It is recommended that the reagent be titrated for optimal performance for each application.
Application Notes:	Additional reported application (for the relevant formats) include: immunohistochemical staining of frozen tissue sections <sup>4,5,8</sup> , immunofluorescent staining <sup>6</sup> , and Western blotting3.
Application References:	<ol> <li>Kan EA, <i>et al.</i> 1983. <i>J. Immunol.</i> 131:536.</li> <li>Wood GS, <i>et al.</i> 1985. <i>Am. J. Pathol.</i> 120:371.</li> <li>Van Dongen JJM, <i>et al.</i> 1988. <i>Blood</i> 71:603. (WB)</li> <li>Haringman JJ, <i>et al.</i> 2005. <i>Arthritis Res. Ther.</i> 7:R862. (IHC)</li> <li>Carbone A, <i>et al.</i> 1999. <i>Blood</i> 93:2319. (IHC)</li> <li>Goval JJ, <i>et al.</i> 2006. <i>J. Histochem. Cytochem.</i> 54:75. (IF)</li> <li>Rutjens E, <i>et al.</i> 2007. <i>J. Immunol.</i> 178:1702.</li> <li>Kap Y, <i>et al.</i> 2009. <i>J. Histochem. Cytochem.</i> 57:1159. (IHC)</li> <li>Yoshino N, <i>et al.</i> 2000. <i>Exp. Anim. (Tokyo)</i> 49:97. (FC)</li> </ol>
Description:	CD3 $\epsilon$ is a 20 kD chain of the CD3/T-cell receptor (TCR) complex, which is composed of two CD3 $\epsilon$ , one CD3 $\gamma$ , one CD3 $\delta$ , one CD3 $\zeta$ (CD247), and a T-cell

·	composed of two CD3 $\epsilon$ , one CD3 $\gamma$ , one CD3 $\delta$ , one CD3 $\zeta$ (CD247), and a T-cell receptor ( $\alpha/\beta$ or $\gamma/\delta$ ) heterodimer. It is found on all mature T cells, NK T cells, and some thymocytes. CD3, also known as T3, is a member of the immunoglobulin superfamily that plays a role in antigen recognition, signal transduction, and T cell activation.
Antigon	1 Barclay N et al 1993 The Leucocyte FactsBook Academic Press San Diego

Antigen	1. Barclay N, <i>et al.</i> 1993. The Leucocyte FactsBook. Academic Press. San Diego.
<b>References:</b>	2. Beverly P, <i>et al.</i> 1981. <i>Eur. J. Immunol.</i> 11:329.
	3. Lanier L, <i>et al.</i> 1986. <i>J. Immunol.</i> 137:2501.

For research use only. Not for diagnostic use. Not for resale. Sony Biotechnology Inc. will not be held responsible for patent infringement or other violations that may occur with the use of our products. Sony Biotechnology Inc. 1730 North First Street, San Jose, CA 95112 www.sonybiotechnology.com