

**PE/Cy7 anti-human CD3**

**Catalog # / Size:** 2324075 / 25 tests  
2324080 / 100 tests

**Clone:** SK7


**Isotype:** Mouse IgG1,  $\kappa$

**Reactivity:** Human

**Preparation:** The antibody was purified by affinity chromatography, and conjugated with PE/Cy7 under optimal conditions. The solution is free of unconjugated PE/Cy7 and unconjugated antibody.

**Formulation:** Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA (origin USA).

**Concentration:** Lot-specific



Human peripheral blood lymphocytes stained with SK7 PE/Cy7

**Applications:**

**Applications:** Flow Cytometry

**Recommended Usage:** Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. **Test size products are transitioning from 20 microL to 5 microL per test.** Please check your vial or your CoA to find the suggested use of this reagent per million cells in 100 microL staining volume or per 100 microL of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

**Application Notes:** Additional reported application (for the relevant formats) include: immunohistochemical staining of frozen tissue sections<sup>4,5,8</sup>, immunofluorescent staining<sup>6</sup>, and Western blotting<sup>3</sup>.

- Application References:**
1. Kan EA, *et al.* 1983. *J. Immunol.* 131:536.
  2. Wood GS, *et al.* 1985. *Am. J. Pathol.* 120:371.
  3. Van Dongen JJM, *et al.* 1988. *Blood* 71:603. (WB)
  4. Haringman JJ, *et al.* 2005. *Arthritis Res. Ther.* 7:R862. (IHC)
  5. Carbone A, *et al.* 1999. *Blood* 93:2319. (IHC)
  6. Goval JJ, *et al.* 2006. *J. Histochem. Cytochem.* 54:75. (IF)
  7. Rutjens E, *et al.* 2007. *J. Immunol.* 178:1702.
  8. Kap Y, *et al.* 2009. *J. Histochem. Cytochem.* 57:1159. (IHC)
  9. Yoshino N, *et al.* 2000. *Exp. Anim. (Tokyo)* 49:97. (FC)

**Description:** CD3 $\epsilon$  is a 20 kD chain of the CD3/T-cell receptor (TCR) complex, which is composed of two CD3 $\epsilon$ , one CD3 $\gamma$ , one CD3 $\delta$ , one CD3 $\zeta$  (CD247), and a T-cell receptor ( $\alpha/\beta$  or  $\gamma/\delta$ ) heterodimer. It is found on all mature T cells, NK T cells, and some thymocytes. CD3, also known as T3, is a member of the immunoglobulin superfamily that plays a role in antigen recognition, signal transduction, and T cell activation.

**Antigen References:**

1. Barclay N, *et al.* 1993. *The Leucocyte FactsBook*. Academic Press. San Diego.
2. Beverly P, *et al.* 1981. *Eur. J. Immunol.* 11:329.
3. Lanier L, *et al.* 1986. *J. Immunol.* 137:2501.