


Brilliant Violet 421™ anti-human CD273 (B7-DC, PD-L2)

Catalog # / 2327595 / 25 tests
Size: 2327600 / 100 tests
Clone: MIH18
Isotype: Mouse IgG1, κ
Immunogen: Human B7-DC transfected cells
Reactivity: Human
Preparation: The antibody was purified by affinity chromatography and conjugated with Brilliant Violet 421™ under optimal conditions. The solution is free of unconjugated Brilliant Violet 421™ and unconjugated antibody.
Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and BSA (origin USA).
Workshop Number: HCDM listed
Concentration: Lot-specific



Human monocyte-derived dendritic cells were stained with CD273 (clone MIH18) Brilliant Violet 421™ (filled histogram) or mouse IgG1, κ isotype control (open histogram).

Applications:

Applications: Flow Cytometry

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is 5 µl per million cells or 5 µl per 100 µl of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

Brilliant Violet 421™ excites at 405 nm and emits at 421 nm. The standard bandpass filter 450/50 nm is recommended for detection. Brilliant Violet 421™ is a trademark of Sirigen Group Ltd.

This product is subject to proprietary rights of Sirigen Inc. and is made and sold under license from Sirigen Inc. The purchase of this product conveys to the buyer a non-transferable right to use the purchased product for research purposes only. This product may not be resold or incorporated in any manner into another product for resale. Any use for therapeutics or diagnostics is strictly prohibited. This product is covered by U.S. Patent(s), pending patent applications and foreign equivalents.

Application Notes: Additional reported applications (for the relevant formats) include: blocking^{4,5}, and immunohistochemistry in frozen sections² and paraffin-embedded formalin-fixed sections⁶.

Application References:
 1. Carreno BM, *et al.* 2002. *Annu. Rev. Immunol.* 20:29.
 2. Ohigashi Y, *et al.* 2005. *Clin. Cancer. Res.* 8:2947.

Description: CD273, known as B7-DC, is also called programmed death ligand 2 (PDL2). This ligand is a 25 kD type I transmembrane protein and a member of B7 family within the immunoglobulin receptor superfamily and is expressed on a subset of dendritic cells, liver and a small subset of macrophages as well as a few transformed cell lines. CD273 has been reported to be stimulatory on dendritic cells when cross-linked and to inhibit T cell activation upon engaging the PD-1 receptor. CD273 has also been reported to bind to an alternative receptor and to mediate T cell activation through such non-PD1 mediated interactions. Clone MIH18 is reported to block PDL2.

Antigen
References:

1. Carreno BM, *et al.* 2002. *Annu. Rev. Immunol.* 20:29.
2. Ohigashi Y, *et al.* 2005. *Clin. Cancer. Res.* 8:2947.