## **Product Data Sheet**

## PE/Cy7 anti-human CD169 (Sialoadhesin, Siglec-1)

Catalog # / 2330065 / 25 tests

Size: 2330070 / 100 tests

Clone: 7-239

Isotype: Mouse IgG1, ĸ

Human Rhinovirus (HRV14) infected, Immunogen:

monocyte derived-DCs

Reactivity: Human

Preparation: The antibody was purified by affinity


chromatography and conjugated with PE/Cy7 under optimal conditions. The solution is free of unconjugated PE/Cy7 and unconjugated antibody.

Phosphate-buffered solution, pH 7.2, Formulation:

containing 0.09% sodium azide and

0.2% (w/v) BSA (origin USA).

Concentration: Lot-specific



IFN- $\gamma$  and TNF- $\alpha$ -stimulated human monocytes (day-3) were

stained with CD169

(Sialoadhesin, Siglec-1, clone 7-239) PE/Cy7 (filled histogram) or Mouse IgG1, κ PE/Cy7 isotype control (open histogram).

## **Applications:**

Applications: Flow Cytometry

Recommended

Usage:

Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is 5 μl per million cells or 5 μl per 100 μl of whole blood. It is recommended that the reagent be titrated for optimal

performance for each application.

**Application** 

Notes:

Additional reported applications (for the relevant formats) include: Western

blotting and inhibition of erythrocyte-rosetting with cells expressing

CD169.

Application

1. Xiong YS, et al. 2009. Clin. Biochem. 42:1057.

References: 2. Varki A, et al. 2009. Glycoconj J. 26:231.

3. Rempel H, et al. 2008. PLoS One. 3:e1967.

4. Crocker PR, et al. 2001. T

**Description:** 

CD169, also known as Siglec-1 and Sialoadhesin (Sn), is a 210 kD type I single membrane-spanning glycoprotein. It is the largest member of the Siglec family, consisting of 1709 amino acids and belonging to the immunoglobulin superfamily. CD169 is expressed by macrophages and dendritic cells. By its affinity to  $\alpha 2,3$ -linked sialic acid, it is involved in macrophage binding to different cell types such as granulocytes,

monocytes, NK, B and T cells. Several CD169 counter receptors, such as CD227 on human breast cancer cells, CD43 on T cells and CD206 on

macrophages, have been reported.

**Antigen** References: 1. Xiong YS, et al. 2009. Clin. Biochem. 42:1057.

2. Varki A, et al. 2009. Glycoconj J. 26:231.

3. Rempel H, et al. 2008. PLoS One. 3:e1967.

4. Crocker PR, et al. 2001. Trends Immunol. 22:337.

5. Hartnell A, et al. 2001. Blood 97:288.

