

**PE anti-human CD360 (IL-21R)**

**Catalog # / Size:** 2339025 / 25 tests  
2339030 / 100 tests

**Clone:** 2G1-K12

**Isotype:** Mouse IgG1, κ


**Immunogen:** IL-21R transfected Ba/F3 cells

**Reactivity:** Human

**Preparation:** The antibody was purified by affinity chromatography, and conjugated with PE under optimal conditions. The solution is free of unconjugated PE and unconjugated antibody.

**Formulation:** Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA (origin USA).


**Concentration:** Lot-specific



**Applications:**

**Applications:** Flow Cytometry

**Recommended Usage:** Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. **Test size products are transitioning from 20 microL to 5 microL per test.** Please check your vial or your CoA to find the suggested use of this reagent per million cells in 100 microL staining volume or per 100 microL of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.



Human peripheral blood lymphocytes were cultured overnight with (bottom) or without (top) recombinant human IL-21, then surface stained with 2G1-K12 PE and CD19 APC

**Application References:** 1. Rodrigues-Bayona B, *et al.* 2012. *J. Immunol.* 188:1578. [PubMed](#)

**Description:** The human interleukin 21 receptor (IL-21R), is a single pass type I membrane protein and a member of the type I cytokine receptor family. Of the type I cytokine receptors, IL-21R exhibits the greatest extracellular homology to the IL-2R β subunit, i.e., contains one copy of the WSXWS-containing cytokine-binding domain. Intracellular domains of IL-21R include the Box 1 and Box 2 elements which are similar to the IL-9R intracellular region. Upon binding IL-21, the IL-21R forms a heterodimer with the common gamma subunit (CD132) and induces Jak/Stat signaling. IL-21R is expressed on B cells and at various levels on NK and T cells. IL-21 is a potent immunomodulatory cytokine mainly produced by NKT and CD4 T-cells (particularly the inflammatory Th17 subset) and has pleiotropic effects on both innate and adaptive immune responses. These actions include

positive effects such as enhanced proliferation of natural killer (NK) cells and cytotoxic T cells that can destroy virally infected or cancerous cells and direct inhibitory effects on the antigen-presenting function of dendritic cells. It can also be proapoptotic for B cells and NK cells. Recent studies have shown that IL-21 is also an autocrine cytokine that potently induces Th17 differentiation and suppresses Foxp3 expression, and serves as a target for treating inflammatory diseases.

**Antigen  
References:**

1. Parish-Novak J, *et al.* 2000. *Nature* 408:57.
2. Ozaki K, *et al.* 2000. *Proc Natl. Acad. Sci. USA.* 97:11439.
3. Dumoutier L, *et al.* 2000. *Proc Natl. Acad. Sci. USA.* 97:10144.