

Brilliant Violet 510™ anti-human IgD

Catalog # / Size: 2341100 / 100 tests
2341095 / 25 tests

Clone: IA6-2

Isotype: Mouse IgG2a, κ


Immunogen: Human IgD

Reactivity: Human

Preparation: The antibody was purified by affinity chromatography and conjugated with Brilliant Violet 510™ under optimal conditions. The solution is free of unconjugated Brilliant Violet 510™ and unconjugated antibody.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and BSA (origin USA).

Concentration: Lot-specific



Human peripheral blood lymphocytes were stained with IgD (clone IA6-2) Brilliant Violet 510™ and CD19 APC.

Applications:

Applications: Flow Cytometry

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is ≤5 microL per million cells or 5 microL per 100 microL of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

Brilliant Violet 510™ excites at 405 nm and emits at 510 nm. The bandpass filter 510/50 nm is recommended for detection, although filter optimization may be required depending on other fluorophores used. **Be sure to verify that your cytometer configuration and software setup are appropriate for detecting this channel.** Refer to your instrument manual or manufacturer for support. Brilliant Violet 510™ is a trademark of Sirigen Group Ltd.

This product is subject to proprietary rights of Sirigen Inc. and is made and sold under license from Sirigen Inc. The purchase of this product conveys to the buyer a non-transferable right to use the purchased product for research purposes only. This product may not be resold or incorporated in any manner into another product for resale. Any use for therapeutics or diagnostics is strictly prohibited. This product is covered by U.S. Patent(s), pending patent applications and foreign equivalents.

Application Notes: Additional reported applications (for the relevant formats) include: immunohistochemical staining of paraformaldehyde fixed frozen sections.⁴

- Application References:**
1. Chen K, *et al.* 2009. *Nat. Immunol.* 10:889.
 2. Lee CH, *et al.* 2005. *J. Exp. Med.* 203:63.
 3. Sutter JA, *et al.* 2008. *Clin. Immunol.* 126:282.
 4. Li H and Pauza CD. 2015. *Eur. J. Immunol.* 45:298. (IHC)

Description: IgD, a member of the immunoglobulin (Ig) family, is expressed in naïve B cells. It has 3 Ig-like domains and exists in a transmembrane and a soluble form. In

general, IgD is not secreted and usually its expression is lost after the Ig isotype switch. After antigen binding, IgD signals through the CD79a/CD79b (Ig α /Ig β) heterodimer, resulting in the activation of the B cell.

**Antigen
References:**

1. Geisberger R, *et al.* 2006. *Immunology* 118:429.
2. Weller S, *et al.* 2005. *Eur. J. Immunol.* 35:2789.
3. Brandtzaeg P and Johansen FE. 2005. *Immunol. Rev.* 206:32.