

PerCP/Cy5.5 anti-human CD134 (OX40)

Catalog # / Size: 2350045 / 25 tests
2350050 / 100 tests

Clone: Ber-ACT35 (ACT35)

Isotype: Mouse IgG1, κ


Immunogen: HTLV 1-transformed HUT 102 cells

Reactivity: Human

Preparation: The antibody was purified by affinity chromatography and conjugated with PerCP/Cy5.5 under optimal conditions. The solution is free of unconjugated PerCP/Cy5.5 and unconjugated antibody.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA (origin USA).

Concentration: Lot-specific




Human peripheral blood lymphocytes were activated for 3 days with PHA, and then stained with CD3 FITC and OX-40 (clone Ber-ACT35) PerCP/Cy5.5 (top) or mouse IgG1, κ PerCP/Cy5.5 isotype control (bottom).

Applications:

Applications: Flow Cytometry

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is 5 microL per million cells or 5 microL per 100 microL of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.



* PerCP/Cy5.5 has a maximum absorption of 482 nm and a maximum emission of 690 nm.

Application Notes: Additional reported applications (for the relevant formats) include: Western blotting¹, immunoprecipitation¹, immunohistochemical staining^{2,3} of paraffin embedded⁷ and frozen tissue sections, ELISA⁴, and functional assay⁵. The LEAF™ or Ultra-LEAF™ purified antibody is recommended for functional assays (contact our [custom solutions team](#)).

Application References:

1. Latza U, *et al.* 1994. *Eur. J. Immunol.* 24:677. (WB, IP)
2. Durkop H, *et al.* 1995. *Brit. J. Haematol.* 91:927. (IHC)
3. Durkop H, *et al.* 1997. *Brit. J. Haematol.* 98:863. (IHC)
4. Willett B, *et al.* 2007. *J. Virol.* 81:9665. (ELISA)

5. Li M and Zhang Y. *et al.* 2005. *Cell. Mol. Immunol.* 2:467. (FA)
 6. Gloviczki ML, *et al.* 2012. *Clin. J. Am. Soc. Nephrol.* 8:546. [PubMed](#)
 7. Domingos PL, *et al.* 2012. *An. Bras. Dermatol.* 87:851. (IHC)
-

Description: CD134, also known as OX40 and TNFRSF4, is a 50 kD type I transmembrane glycoprotein. It is a member of the TNF receptor family. OX40 is expressed on activated T lymphocytes including Th1, Th2, Th17, and Treg cells. The interaction of OX40 with OX40L results in B cell proliferation and antibody secretion, regulation of primary T cell expansion, and T cell survival. OX40 influences the size of the T cell memory pool and regulation of CD4⁺ T cell tolerance.

- Antigen**
- References:**
1. Smith CA, *et al.* 1994. *Cell.* 76:959.
 2. Chen AL, *et al.* 1999. *Immunity.* 11:689.
 3. Croft M. 2010. *Annu. Rev. Immunol.* 28:57.
 4. Ruby CE, *et al.* 2009. *J. Immunol.* 183:5079