

**APC anti-human CD325 (N-Cadherin)**

**Catalog # /** 2354040 / 100 tests  
**Size:** 2354035 / 25 tests

**Clone:** 8C11

**Isotype:** Mouse IgG1,  $\kappa$


**Immunogen:** Recombinant human N-cadherin extracellular domain

**Reactivity:** Human

**Preparation:** The antibody was purified by affinity chromatography and conjugated with APC under optimal conditions. The solution is free of unconjugated APC and unconjugated antibody.

**Formulation:** Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA (origin USA).

**Concentration:** Lot-specific



Human acute myeloid leukemia cell line (KG1) was stained with CD325 (clone 8C11) APC (filled histogram) or mouse IgG1,  $\kappa$  APC isotype control (open histogram).

**Applications:**

**Applications:** Flow Cytometry

**Recommended Usage:** Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. **Test size products are transitioning from 20 microl to 5 microl per test.** Please check your vial or your CoA to find the suggested use of this reagent per million cells in 100 microl staining volume or per 100 microl of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

**Application Notes:** The mAb 8C11 recognizes the amino acids 92-593 of CD325, located between the extracellular cadherin structural domain (EC) 3 and 4. Additional reported applications (for the relevant formats) include: immunofluorescence<sup>1,3,6</sup>, motility inhibition of N-cadherin-expressing cells<sup>2</sup>, and Western blot<sup>2,4</sup>.

**Application References:**

1. Navarro P, et al. 1998. *J. Cell Biol.* 140:1475. (IF)
2. Kim JB. 2000. *J. Cell Biol.* 151:1193. (Block, WB)
3. Puch S, et al. 2001. *J. Cell. Sci.* 114:1567. (IF)
4. Wahl JK. 3rd, et al. 2003. *J. Biol. Chem.* 278:17269. (WB)
5. Wein F, et al. 2010. *Stem. Cell Res.* 4:129. (FC)
6. Jaggi M, et al. 2002. *Cell. Commun. Adhes.* 9:103. (IF)

**Description:** CD325 (N-cadherin) is a 130 kD, single pass transmembrane protein. Its extracellular region consists of five EC domains and has one cytoplasmic domain. N-cadherin is involved in organogenesis and maintenance of organ architecture by contributing to the sorting of heterogeneous cell types and in the cell adhesion needed to form tissues. N-cadherin is expressed by stem cells, myeloblasts, endothelial cells, and fibroblasts, and also is expressed in neural and muscle tissues and some types of carcinoma cells. CD325 associates with the cytoskeleton through catenin proteins.

**Antigen References:**

1. Colomiere M, et al. 2009. *Brit. J. Cancer* 100:134.
2. Yan W, et al. 2010. *J. Biol. Chem.* 285:14042.
3. Mosnier JF, et al. 2009. *Mod. Pathol.* 22:182.
4. Gao L, et al. 201

