
## **Product Data Sheet**

## PE/Cy7 anti-human CD276 (B7-H3)

Catalog # / Size:	2355040 / 100 tests 2355035 / 25 tests	
Clone:	MIH42	
Isotype:	Mouse lgG1, κ	nber
Immunogen:	Human B7-H3	Relative Cell Number
<b>Reactivity:</b>	Human	he Ce
Preparation:	The antibody was purified by affinity chromatography and conjugated with PE/Cy7 under optimal conditions. The solution is free of unconjugated PE/Cy7 and unconjugated antibody.	Relat
Formulation:	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA (origin USA).	Huce
Concentration:	Lot-specific	(c hi:



Human B7-H3 transfected P815 cells were stained with CD276 (clone MIH42) PE/Cy7 (filled histogram) or mouse IgG1, κ PE/Cy7 isotype control (open histogram).

## **Applications:**

Applications: Flow Cytometry

**Recommended** Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is 5 microL per million cells or 5 microL per 100 microL of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

**Description:** B7-H3, assigned as CD276, is a type I transmembrane protein and shares 20-27% amino acid identity with other B7 family members. Human B7-H3 has a single extracellular variable-type immunoglobulin (IgV)-IgC domain, a signature intracellular domain, and an additional isoform, known as 4Ig-B7-H3, containing nearly exact tandem duplication of the IgV-IgC domain and most likely caused by exon duplication. B7-H3 mRNA is broadly expressed in normal tissues whereas its protein expression is relatively rare. The expression of B7-H3 is induced on T cells, natural killer (NK) cells, and antigen-presenting cells (APCs), including dendritic cells (DCs) and macrophages. It can be upregulated during the maturation from monocytes to DCs, or during the interaction between DCs and regulatory T cells. B7-H3 has been shown to be a co-stimulatory molecule that inhibits T-cell responses. B7-H3 has also been identified to bind TLT-2 involved in the intracellular signaling pathway.

For research use only. Not for diagnostic use. Not for resale. Sony Biotechnology Inc. will not be held responsible for patent infringement or other violations that may occur with the use of our products. Sony Biotechnology Inc. 1730 North First Street, San Jose, CA 95112 www.sonybiotechnology.com