## **Product Data Sheet**

## PE/Cy7 anti-human TCR Vα7.2

**Catalog # / Size:** 2358560 / 100 tests

2358555 / 25 tests

Clone: 3C10

**Isotype:** Mouse IgG1, κ

Immunogen: Recombinant TCR

Reactivity: Human

**Preparation:** The antibody was purified by affinity

chromatography and conjugated with PE/Cy7 under optimal conditions. The solution is free of unconjugated PE/Cy7


and unconjugated antibody.

Formulation: Phosphate-buffered solution, pH 7.2,

containing 0.09% sodium azide and

0.2% (w/v) BSA (origin USA).

Concentration: Lot-specific



Human peripheral blood lymphocytes were stained with CD3 FITC and TCR V $\alpha$ 7.2 (clone 3C10, top) PE/Cy7 or mouse IgG1,  $\kappa$ PE/Cy7isotype control (bottom).

## **Applications:**

**Applications:** Flow Cytometry

Recommended

**Usage:** 

Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is 5 microL per million cells or 5 microL per 100 microL of whole blood. It is recommended that the reagent be titrated for optimal performance for

each application.

Application Notes:

Associated with an anti-CD161 or - IL18R $\alpha$  staining, the 3C10 antibody allows unequivocal identification of MAIT cells. Importantly, the V $\alpha$ 7.2 segment can also be used by conventional T cells. Therefore, the 3C10 also stains a subset of conventional CD4 and CD8 T

cells.

**Application** 

1. Martin E, et al. 2009. PLoS Biol. 7:525.

**References:** 

2. Wakao H, et al. 2013. Cell Stem Cell 12:1. PubMed

**Description:** 

The 3C10 antibody recognizes the V $\alpha$ 7.2 T cell antigen receptor (TCR)  $\alpha$ -chain segment which, joined with the J $\alpha$ 33 segment, constitutes an invariant TCR that is a characteristic of the mucosal-associated invariant T cells (MAIT cells). MAIT cells are restricted by a nonpolymorphic class Ib major histocompatibility complex (MHC) molecule, MHC-related molecule 1 (MR1). MAIT cells are present in human blood (1-8% of T cells), mesenteric lymph nodes, liver, and intestinal mucosa. MAIT cells play a role in detecting and fighting off microbial infections.

lgG1, k PE/Cy7 isotype control

103

10<sup>2</sup>

104

CD3 FITC

105

Antigen References:	1. Le Bourhis L, <i>et al.</i> 2010. <i>Nat. Immunol.</i> 11:701.