Product Data Sheet

APC/Cy7 anti-human TCR Vα7.2

Catalog # / Size: 2358565 / 25 tests

2358570 / 100 tests

Clone:

Isotype: Mouse IgG1, κ

Recombinant TCR Immunogen:

Reactivity: Human

Preparation: The antibody was purified by affinity

chromatography and conjugated with APC/Cy7 under optimal conditions. The solution is free of unconjugated APC/Cy7


and unconjugated antibody.

Formulation: Phosphate-buffered solution, pH 7.2,

containing 0.09% sodium azide and

0.2% (w/v) BSA (origin USA).

Concentration: Lot-specific



Human peripheral blood lymphocytes were stained with CD3 FITC and TCR $V\alpha7.2$ (clone 3C10. top) APC/Cv7 or mouse IaG1, k APC/Cy7 isotype control (bottom).

Applications:

Applications: Flow Cytometry

Recommended

Usage:

Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is 5 microL per million cells or 5 microL per 100 microL of whole blood. It is recommended that the reagent be titrated for optimal performance for

each application.

Application Notes:

Associated with an anti-CD161 or -IL18Rα staining, the 3C10 antibody allows unequivocal identification of MAIT cells. Importantly, the $V\alpha7.2$ segment can also be used by conventional T cells. Therefore, the 3C10 also stains a subset of conventional CD4 and CD8 T

cells.

Application

1. Martin E, et al. 2009. PLoS Biol. 7:525.

References:

2. Wakao H, et al. 2013. Cell Stem Cell 12:1. PubMed

Description:

The 3C10 antibody recognizes the V α 7.2 T cell antigen receptor (TCR) α -chain segment which, joined with the J α 33 segment, constitutes an invariant TCR that is a characteristic of the mucosal-associated invariant T cells (MAIT cells). MAIT cells are restricted by a nonpolymorphic class Ib major histocompatibility complex (MHC) molecule, MHC-related molecule 1 (MR1). MAIT cells are present in human blood (1-8% of T cells), mesenteric lymph nodes, liver, and intestinal mucosa. MAIT cells play a role in detecting and fighting off microbial infections.

Mouse IgG1, x APC/Cy7

102

0

104

102 CD3 FITC

Antigen References:	1. Le Bourhis L, <i>et al.</i> 2010. <i>Nat. Immunol.</i> 11:701.