

Brilliant Violet 785™ anti-human TCR Vα7.2

Catalog # / Size: 2358605 / 25 tests
2358610 / 100 tests

Clone: 3C10

Isotype: Mouse IgG1, κ


Immunogen: Recombinant TCR

Reactivity: Human

Preparation: The antibody was purified by affinity chromatography and conjugated with Brilliant Violet 785™ under optimal conditions. The solution is free of unconjugated Brilliant Violet 785™ and unconjugated antibody.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and BSA (origin USA).

Concentration: 0.2




Human peripheral blood lymphocytes were stained with CD3 APC and TCR Vα7.2 (clone 3C10) Brilliant Violet 785™ (top), or mouse IgG1, κ Brilliant Violet 785™ isotype control (bottom).

Applications:

Applications: Flow Cytometry

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is ≤5 microL per million cells or 5 microL per 100 microL of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.



Brilliant Violet 785™ excites at 405 nm and emits at 785 nm. The bandpass filter 780/60 nm is recommended for detection, although filter optimization may be required depending on other fluorophores used. **Be sure to verify that your cytometer configuration and software setup are appropriate for detecting this channel.** Refer to your instrument manual or manufacturer for support. Brilliant Violet 785™ is a trademark of Sirigen Group Ltd.

Application Notes: Associated with an anti-CD161 or -IL18Rα staining, the 3C10 antibody allows unequivocal identification of MAIT cells. Importantly, the Vα7.2 segment can also be used by conventional T cells. Therefore, the 3C10 also stains a subset of conventional CD4 and CD8 T

cells.

- Application** 1. Martin E, *et al.* 2009. *PLoS Biol.* 7:525.
References: 2. Wakao H, *et al.* 2013. *Cell Stem Cell* 12:1. [PubMed](#)
-

Description: The 3C10 antibody recognizes the V α 7.2 T cell antigen receptor (TCR) α -chain segment which, joined with the J α 33 segment, constitutes an invariant TCR that is a characteristic of the mucosal-associated invariant T cells (MAIT cells). MAIT cells are restricted by a nonpolymorphic class Ib major histocompatibility complex (MHC) molecule, MHC-related molecule 1 (MR1). MAIT cells are present in human blood (1-8% of T cells), mesenteric lymph nodes, liver, and intestinal mucosa. MAIT cells play a role in detecting and fighting off microbial infections.

- Antigen**
References: 1. Le Bourhis L, *et al.* 2010. *Nat. Immunol.* 11:701.