

Pacific Blue™ anti-human CD63

Catalog # / Size: 2365055 / 25 tests
2365060 / 100 tests

Clone: H5C6

Isotype: Mouse IgG1, κ


Immunogen: T cell line HPB-ALL

Reactivity: Human

Preparation: The antibody was purified by affinity chromatography, and conjugated with Pacific Blue™ under optimal conditions. The solution is free of unconjugated Pacific Blue™.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA (origin USA).

Concentration: Lot-specific



Thrombin-activated human peripheral blood platelets were stained with CD63 (clone H5C6) Pacific Blue™ (filled histogram) or mouse IgG1, κ Pacific Blue™ isotype control (open histogram).

Applications:

Applications: Flow Cytometry

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is 5 microL per million cells or 5 microL per 100 microL of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

* Pacific Blue™ has a maximum emission of 455 nm when it is excited at 405 nm. Prior to using Pacific Blue™ conjugate for flow cytometric analysis, please verify your flow cytometer's capability of exciting and detecting the fluorochrome.

Application Notes: Additional reported applications (for the relevant formats) include: Western blotting¹, immunofluorescence², and immunoprecipitation¹.

- Application References:**
1. Hildreth JE, *et al.* 1991. *Blood* 77:121. (IP, WB)
 2. Beatty WL, *et al.* 2006. *J. Cell Sci.* 119:350. (IF)
 3. Liang Y, *et al.* 2014. *J Biol Chem.* 289:32526. [PubMed](#)

Description: CD63 is a 53 kD type III lysosomal glycoprotein also known as LIMP, LAMP-3, gp55, and melanoma-associated antigen (ME491). CD63 is a member of the tetraspan transmembrane superfamily (TM4SF) protein expressed on activated platelets, monocytes/macrophages, endothelium, fibroblasts, osteoclasts, and smooth muscle cells. CD63 may be involved in platelet activation and is thought to function as a transmembrane adaptor protein. CD63 has been shown to associate with CD9, CD81, VLA-3, and VLA-6.

- Antigen References:**
1. Azorsa DO, *et al.* 1991. *Blood* 78:280.
 2. Kishimoto T, *et al.* Eds. 1997. *Leukocyte Typing V1.* Oxford University Press New York.
 3. Hildreth JE, *et al.* 1991. *Blood* 77:121.
 4. Anzai N, *et*

