Product Data Sheet

Brilliant Violet 711™ anti-human CD63

Catalog # / 2365205 / 25 tests

Size: 2365210 / 100 tests

Clone: H5C6

Isotype: Mouse IgG1, κ

Immunogen: T cell line HPB-ALL

Reactivity: Human, Non-human primate, Other

Preparation: The antibody was purified by affinity

chromatography and conjugated with Brilliant Violet 711^{TM} under optimal conditions. The solution is free of unconjugated Brilliant Violet 711^{TM}

and unconjugated antibody.

Formulation: Phosphate-buffered solution, pH 7.2,

containing 0.09% sodium azide and

BSA (origin USA).

Workshop Number: **HCDM** listed

Concentration: Lot-specific

Thrombin-activated human peripheral blood platelets were stained with CD63 (clone H5C6) Brilliant Violet 711™ (filled histogram) or Mouse IgG1, κ Brilliant Violet 711™ isotype control (open histogram).

Applications:

Applications: Flow Cytometry

Human peripheral blood monocytes were stained with HLA-DR FITC and Brilliant Violet 421™ anti-human CD14 (clone 63D3) (left) or Brilliant Violet 421™ mouse IgG1, κ isotype control (right).

Recommended Usage:

Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is 5 μ l per million cells in 100 μ l staining volume or 5 μ l per 100 μ l of whole blood.

Brilliant Violet 711™ excites at 405 nm and emits at 711 nm. The bandpass filter 710/50 nm is recommended for detection, although filter optimization may be required depending on other fluorophores used. Be sure to verify that your cytometer configuration and software setup are appropriate for detecting this channel. Refer to your instrument manual or manufacturer for support. Brilliant Violet 711™ is a trademark of Sirigen Group Ltd.

This product is subject to proprietary rights of Sirigen Inc. and is made and sold under license from Sirigen Inc. The purchase of this product conveys to the buyer a non-transferable right to use the purchased product for research purposes only. This product may not be resold or incorporated in any manner into another product for resale. Any use for therapeutics or diagnostics is strictly prohibited. This product is covered by U.S. Patent(s), pending patent applications and foreign equivalents.

Application Notes:

Additional reported applications (for the relevant formats) include:

Western blotting¹,

 $immun of luorescence^2, \ and \\ immun oprecipitation^1.$

Application References:

- 1. Hildreth JE, et al. 1991. Blood 77:121. (IP, WB)
- 2. Beatty WL, et al. 2006. J. Cell Sci. 119:350. (IF)

Description:

CD63 is a 53 kD type III lysosomal glycoprotein also known as LIMP, LAMP-3, gp55, and melanoma-associated antigen (ME491). CD63 is a member of the tetraspan transmembrane superfamily (TM4SF) protein expressed on activated platelets, monocytes/macrophages, endothelium, fibroblasts, osteoclasts, and smooth muscle cells. CD63 may be involved in platelet activation and is thought to function as a transmembrane adaptor protein. CD63 has been shown to associate with CD9, CD81, VLA-3, and VLA-6.

Antigen References:

- 1. Azorsa DO, et al. 1991. Blood 78:280.
- 2. Kishimoto T, et al. Eds. 1997. Leukocyte Typing V1. Oxford University Press New York.
- 3. Hildreth JE, et al. 1991. Blood 77:121.
- 4. Anzai N, et al. 2002. Blood 99:4413.