

Brilliant Violet 421™ anti-human CD54

Catalog # / Size: 2365655 / 25 tests
2365660 / 100 tests

Clone: HA58

Isotype: Mouse IgG1, κ

Immunogen: Colonic cancer BM314 cells


Reactivity: Human

Preparation: The antibody was purified by affinity chromatography and conjugated with Brilliant Violet 421™ under optimal conditions.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and BSA (origin USA)

Workshop Number: HCDM listed

Concentration: Lot-specific



Human peripheral blood lymphocytes were stained with CD54 (clone HA58) Brilliant Violet 421™ (filled histogram) or mouse IgG1, κ Brilliant Violet 421™ isotype control (open histogram).

Applications:

Applications: Flow Cytometry

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is 5 μL per million cells in 100 μL staining volume or 5 μL per 100 μL of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

Brilliant Violet 421™ excites at 405 nm and emits at 421 nm. The standard bandpass filter 450/50 nm is recommended for detection. Brilliant Violet 421™ is a trademark of Sirigen Group Ltd.

This product is subject to proprietary rights of Sirigen Inc. and is made and sold under license from Sirigen Inc. The purchase of this product conveys to the buyer a non-transferable right to use the purchased product for research purposes only. This product may not be resold or incorporated in any manner into another product for resale. Any use for therapeutics or diagnostics is strictly prohibited. This product is covered by U.S. Patent(s), pending patent applications and foreign equivalents.

Application Notes: Clone HA58 recognizes an epitope located in the extracellular D1 domain of CD54.³

- Application References:**
1. Tsujisaki M, *et al.* 1991. *Clin. Exp. Immunol.* 85:3.
 2. Kanwar JR, *et al.* 2003. *Cancer Gene Ther.* 10:468.
 3. Kohka H, *et al.* 1998. *J. Leukoc. Biol.* 64:519.

Description: CD54 is a 85-110 kD type I transmembrane protein also known as ICAM-1. It is expressed on activated endothelial cells, high endothelial venules, T and B cells, monocytes/macrophages, granulocytes, and dendritic cells. The expression of ICAM-1 can be released from the cell surface. CD54 plays a role in cellular adhesion and is involved in inflammation and leukocyte extravasation. CD54 has also been shown to be the major cellular receptor for rhinovirus. ICAM-1 binds to CD11a/CD18 (LFA-1), CD11b/CD18 (Mac-1), CD11c/CD18 (p150, 95) as well as hyaluronan and fibrinogen.

Antigen
References:

1. Voraberger G, *et al.* 1991. *J. Immunol.* 147:2777.
2. Staunton DE, *et al.* 1988. *Cell* 52:925.
3. Greve JM, *et al.* 1989. *Cell* 56:839.