Product Data Sheet

PE anti-human CD183 (CXCR3)

Catalog # / Size: 2368530 / 100 tests

2368525 / 25 tests

Clone: G025H7

Isotype: Mouse IgG1, κ

Immunogen: Human CXCR3 transfectants

Reactivity: Human

Preparation: The antibody was purified by affinity

chromatography and conjugated with PE under optimal conditions. The solution is free of unconjugated PE and


unconjugated antibody.

Formulation: Phosphate-buffered solution, pH 7.2,

containing 0.09% sodium azide and

0.2% (w/v) BSA (origin USA).

Concentration: Lot-specific



Human peripheral blood lymphocytes were stained with CD3 APC and CXCR3 (clone G025H7) PE (top plot) or mouse IgG1 PE isotype control (bottom plot).

Applications:

Applications: Flow Cytometry


Recommended

Usage:

Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis.

Test size products are transitioning from 20 microL to 5 microL per test.

Please check your vial or your CoA to find the suggested use of this reagent per million cells in 100 microL staining volume or per 100 microL of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.



Description:

Human CXCR3, also known as GPR9, is a chemokine receptor that binds CXCL9, CXCL10, and CXCL11. It is a 38 kD seven-pass transmembrane receptor coupled to G-protein. CXCR3 is highly expressed by T cells (Th1), natural killer cells (NK cells), dendritic cells, mast cells, alveolar macrophages, eosinophils, and human airway epithelial cells. CXCR3 is important for effector lymphocyte recruitment into inflamed tissue in various inflammatory and autoimmune diseases, such as chronically inflamed liver, Crohn's disease, rheumatoid arthritis, multiple sclerosis, and inflammatory skin diseases.

Antigen References:

- 1. Loetscher M, et al. 1996. J. Exp. Med. 184:963.
- 2. Cole KE, et al. 1998. J. Exp. Med. 187:2009.
- 3. Aksov MO, et al. 2006. Am. J. Physiol. Lung Cell Mol. Physiol. 290:L909.
- 4. Curbi