


Purified anti-human CD303 (BDCA-2)


Catalog # / Size: 2371010 / 100 µg
Clone: 201A
Isotype: Mouse IgG2a, κ
Reactivity: Human
Preparation: The antibody was purified by affinity chromatography.
Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.
Concentration: 0.5



Human peripheral mononuclear cells were stained with purified CD303 (clone 201A, top) or purified mouse IgG2a, κ isotype control (bottom), followed by anti-mouse IgG PE and CD123 APC. Data shown was gated on the lymphocyte and monocyte populations.

Applications:

Applications: Flow Cytometry
Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is ≤1.0 microg per million cells in 100 microL volume. It is recommended that the reagent be titrated for optimal performance for each application.



Description: CD303, also known as BDCA-2 and CLEC4C, is a 38 kD type II transmembrane glycoprotein. It is a member of the C-type lectin superfamily. CD303 is expressed by plasmacytoid dendritic cells (pDCs) and is involved in cell adhesion, signaling, and antigen capture and processing. Crosslinking of CD303 inhibits the production of IFN-α/β and TLR-9 induced pDCs maturation.

- Antigen References:**
1. Jähn PS, *et al.* 2010. *Cell Immunol.* 265:15.
 2. Graham LM and Brown GD. 2009. *Cytokine* 48:148.
 3. Röck J, *et al.* 2007. *Eur. J. Immunol.* 37:3564.
 4. Dzionek A, *et al.* 20