

Brilliant Violet 510™ anti-human CD303 (BDCA-2)

Catalog # / Size: 2371160 / 100 tests
2371155 / 25 tests

Clone: 201A


Isotype: Mouse IgG2a, κ

Reactivity: Human

Preparation: The antibody was purified by affinity chromatography and conjugated with Brilliant Violet 510™ under optimal conditions. The solution is free of unconjugated Brilliant Violet 510™ and unconjugated antibody.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and BSA (origin USA).

Concentration: Lot-specific



Human peripheral blood lymphocytes were stained with CD123 APC, and CD303 (BDCA-2, clone 201A) Brilliant Violet 510™ (left) or mouse IgG1, κ Brilliant Violet 510™ isotype control (right).

Applications:

Applications: Flow Cytometry

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is 5 µl per million cells or 5 µl per 100 µl of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

Brilliant Violet 510™ excites at 405 nm and emits at 510 nm. The bandpass filter 510/50 nm is recommended for detection, although filter optimization may be required depending on other fluorophores used. Be sure to verify that your cytometer configuration and software setup are appropriate for detecting this channel. Refer to your instrument manual or manufacturer for support. Brilliant Violet 510™ is a trademark of Sirigen Group Ltd.

This product is subject to proprietary rights of Sirigen Inc. and is made and sold under license from Sirigen Inc. The purchase of this product conveys to the buyer a non-transferable right to use the purchased product for research purposes only. This product may not be resold or incorporated in any manner into another product for resale. Any use for therapeutics or diagnostics is strictly prohibited. This product is covered by U.S. Patent(s), pending patent applications and foreign equivalents.

- Application References:**
1. Jähn PS, *et al.* 2010. *Cell Immunol.* 265:15.
 2. Graham LM and Brown GD. 2009. *Cytokine* 48:148.
 3. Röck J, *et al.* 2007. *Eur. J. Immunol.* 37:3564.
 4. Dzionek A, *et al.* 20

Description: CD303, also known as BDCA-2 and CLEC4C, is a 38 kD type II transmembrane glycoprotein. It is a member of the C-type lectin superfamily. CD303 is expressed by plasmacytoid dendritic cells (pDCs) and is involved in cell adhesion, signaling,

and antigen capture and processing. Crosslinking of CD303 inhibits the production of IFN- α/β and TLR-9 induced pDCs maturation.

**Antigen
References:**

1. Jähn PS, *et al.* 2010. *Cell Immunol.* 265:15.
2. Graham LM and Brown GD. 2009. *Cytokine* 48:148.
3. Röck J, *et al.* 2007. *Eur. J. Immunol.* 37:3564.
4. Dzionek A, *et al.* 20