

**PE anti-human CD304 (Neuropilin-1)**

**Catalog # / Size:** 2372520 / 100 tests  
2372515 / 25 tests

**Clone:** 12C2

**Isotype:** Mouse IgG2a, κ


**Immunogen:** CD304-Fc Fusion protein

**Reactivity:** Human

**Preparation:** The antibody was purified by affinity chromatography and conjugated with PE under optimal conditions. The solution is free of unconjugated PE and unconjugated antibody.

**Formulation:** Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA (origin USA).

**Concentration:** Lot-specific




Human peripheral blood mononuclear cells were stained with CD303 APC and CD304 (clone 12C2) PE (top) or mouse IgG2a, κ PE isotype control (bottom). Data shown was gated on the lymphocyte and monocyte populations.

**Applications:**

**Applications:** Flow Cytometry

**Recommended Usage:** Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is 5 microL per million cells or 5 microL per 100 microL of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.



**Description:** CD304, also known as neuropilin-1, BDCA-4 and VEGF165R, is a 140 kD type I transmembrane protein. Its extracellular region contains 2 CUB, 2 FV/FVIII, and one MAM domain; a soluble isoform is produced by alternative mRNA splicing. CD304 is involved in angiogenesis, neural development, and tumor metastasis. It's expressed by plasmacytoid dendritic cells, thymocytes, neurons, endothelium, and a subset of T<sub>FH</sub> cells. CD304 is also expressed in several carcinomas, and a high expression of this molecule in prostate cancer correlates with a poor prognosis.

- Antigen References:**
1. Mizui M and Kikutani H. 2008. *Immunity* 28:302.
  2. Hamerlik P, et al. 2012. *J. Exp. Med.* 209:507.
  3. Karjalainen K, et al. 2011. *Blood* 117:920.
  4. Lepelletier Y, et al. 2007.