

PE/Cy7 anti-human CD21

Catalog # / Size: 2374555 / 25 tests
2374560 / 100 tests

Clone: Bu32

Isotype: Mouse IgG1, κ


Reactivity: Human

Preparation: The antibody was purified by affinity chromatography and conjugated with PE/Cy7 under optimal conditions. The solution is free of unconjugated PE/Cy7 and unconjugated antibody.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA (origin USA).

Workshop Number: V CD21.4, VI CD21.5

Concentration: Lot-specific



Human peripheral blood lymphocytes were stained with CD19 Brilliant Violet 421™ and CD21 (clone Bu32) PE/Cy7 (top) or mouse IgG1, κ PE/Cy7 isotype control (bottom).


Applications:

Applications: Flow Cytometry

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is 5 microL per million cells or 5 microL per 100 microL of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

Application Notes: Additional reported applications (for the relevant formats) include: immunohistochemical staining of acetone-fixed frozen sections⁴.

- Application References:**
1. Björck P, *et al.* 1993. *Eur. J. Immunol.* 23:1771.
 2. Frémeaux-Bacchi V, *et al.* 1996. *Eur. J. Immunol.* 26:1497.
 3. Ling NR, *et al.* 1995. *Clin. Exp. Immunol.* 101:369.
 4. Wang, C, *et al.* 2011. *BMC Immunol.* 12:53. (IHC)



Description: CD21 is a 145 kD transmembrane protein also known as complement C3d receptor (C3dR), complement receptor 2 (CR2), and Epstein-Barr virus receptor. CD21 is expressed on B cells, follicular dendritic cells, subsets of normal thymocytes and T cells, and some epithelial cells. CD21 is the receptor used by Epstein-Barr virus to infect B cells and is also the complement receptor for C3d. CD21 has also been shown to interact with a number of proteins, including CD23, CD19, annexin VI, CD81, iC3b, complement receptor 1 (CR1, CD35), and interferon- α 1 (IFN- α 1).

- Antigen**
- References:**
1. Kishimoto T, Eds. 1997. Leukocyte Typing VI. Garland Publishing Inc.
 2. Moore MD, *et al.* 1987. *Proc. Natl. Acad. Sci. USA* 84:9194.
 3. Szakonyi G, *et al.* 2001. *Science* 292:1725.
 4. Weis JJ, *et*