


Purified anti-human CD27

Catalog # / Size: 2382005 / 100 µg
Clone: M-T271
Isotype: Mouse IgG1, κ
Immunogen: Human T cells from a T-ALL patient.
Reactivity: Human
Preparation: The antibody was purified by affinity chromatography.
Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.
Workshop Number: V 5T CD27.03
Concentration: 0.5



Human peripheral blood lymphocytes were stained with purified CD27 (clone M-T271) (filled histogram) or mouse IgG1 isotype control (open histogram), followed by anti-mouse IgG PE.

Applications:

Applications: Other

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is ≤0.5 microg per million cells in 100 microL volume. It is recommended that the reagent be titrated for optimal performance for each application.

Application Notes: Additional reported applications (for the relevant formats) include: immunohistochemical staining of formalin-fixed paraffin-embedded frozen tissue sections¹, immunofluorescent staining², and ELISA³.

Application References: 1. Ma S, *et al.* 2011. *J. Virol.* 85:165. (IHC)
2. Manzo A, *et al.* 2008. *Arthritis Rheum.* 11:3377. (IF)
3. Kato K, *et al.* 2007. *Exp. Hematol.* 35:434. (ELISA)

Description: CD27 is a 50-55 kD type I membrane protein also known as S152 and T14. It is a lymphocyte-specific member of the TNF-receptor superfamily. CD27 is expressed on medullary thymocytes, virtually all mature T cells, some B cells, and NK cells. CD27 binds to CD70, and plays a role in costimulation of T cell activation and regulation of B cell differentiation and proliferation. The cytoplasmic domains of CD27 have also been shown to interact with TRAF2 and TRAF5 to elicit NF-κB and SAPK/JNK activation.

Antigen References: 1. Knapp W, *et al.* 1989. *Leucocyte Typing IV: White Cell Differentiation Antigens.* Oxford University Press.
2. Schlossman S, *et al.* 1995. *Leucocyte Typing V: White Cell Differentiation Antigens.* Oxford University Press.
3.