FITC anti-human CD27

Catalog # / Size: 2382020 / 100 tests

2382015 / 25 tests

Clone: M-T271

Isotype: Mouse IgG1, κ

Human T cells from a T-ALL patient. Immunogen:

Reactivity: Human

Preparation: The antibody was purified by affinity

chromatography and conjugated with FITC under optimal conditions. The solution is free of unconjugated FITC

and unconjugated antibody.

Formulation: Phosphate-buffered solution, pH 7.2,


containing 0.09% sodium azide and

0.2% (w/v) BSA (origin USA).

Workshop **Number:**

V 5T CD27.03

Concentration: Lot-specific



Human peripheral blood lymphocytes were stained with CD27 (clone M-T271) FITC (filled histogram) or mouse IgG1 FITC isotype control (open histogram).

Applications:

Applications: Flow Cytometry

Recommended Each lot of this antibody is quality control tested by immunofluorescent staining

with flow cytometric analysis. For flow cytometric staining, the suggested use of Usage:

this reagent is 5 microL per million cells or 5 microL per 100 microL of whole blood. It is recommended that the reagent be titrated for optimal performance for

each application.

Application Additional reported applications (for the relevant formats) include:

Notes:

immunohistochemical staining of formalin-fixed paraffin-embedded frozen tissue

sections1, immunofluorescent staining2, and ELISA3.

1. Ma S, et al. 2011. J. Virol. 85:165. (IHC) **Application**

References: 2. Manzo A, et al. 2008. Arthritis Rheum. 11:3377. (IF)

3. Kato K, et al. 2007. Exp. Hematol. 35:434. (ELISA)

CD27 is a 50-55 kD type I membrane protein also known as S152 and T14. It is a **Description:**

> lymphocyte-specific member of the TNF-receptor superfamily. CD27 is expressed on medullary thymocytes, virtually all mature T cells, some B cells, and NK cells. CD27 binds to CD70, and plays a role in costimulation of T cell activation and regulation of B cell differentiation and proliferation. The cytoplasmic domains of CD27 have also been shown to interact with TRAF2 and TRAF5 to elicit NF-кВ and

SAPK/INK activation.

Antigen References: 1. Knapp W, et al. 1989. Leucocyte Typing IV: White Cell Differentiation Antigens.

Oxford University Press.

2. Schlossman S, et al. 1995. Leucocyte Typing V: White Cell Differentiation

Antigens. Oxford University Press.

3.