


Brilliant Violet 510™ anti-human CD27

Catalog # / Size: 2382100 / 100 tests
Clone: M-T271
Isotype: Mouse IgG1, κ
Immunogen: Human T cells from a T-ALL patient.
Reactivity: Human
Preparation: The antibody was purified by affinity chromatography and conjugated with Brilliant Violet 510™ under optimal conditions. The solution is free of unconjugated Brilliant Violet 510™ and unconjugated antibody.
Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and BSA (origin USA).
Workshop Number: V 5T CD27.03
Concentration: Lot-specific



Human peripheral blood lymphocytes were stained with CD27 (clone M-T271) Brilliant Violet 510™ (filled histogram) or mouse IgG1 Brilliant Violet 510™ isotype control (open histogram).

Applications:

Applications: Flow Cytometry
Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is ≤5 microL per million cells or 5 microL per 100 microL of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

Brilliant Violet 510™ excites at 405 nm and emits at 510 nm. The bandpass filter 510/50 nm is recommended for detection, although filter optimization may be required depending on other fluorophores used. **Be sure to verify that your cytometer configuration and software setup are appropriate for detecting this channel.** Refer to your instrument manual or manufacturer for support. Brilliant Violet 510™ is a trademark of Sirigen Group Ltd.

This product is subject to proprietary rights of Sirigen Inc. and is made and sold under license from Sirigen Inc. The purchase of this product conveys to the buyer a non-transferable right to use the purchased product for research purposes only. This product may not be resold or incorporated in any manner into another product for resale. Any use for therapeutics or diagnostics is strictly prohibited. This product is covered by U.S. Patent(s), pending patent applications and foreign equivalents.

Application Notes: Additional reported applications (for the relevant formats) include: immunohistochemical staining of formalin-fixed paraffin-embedded frozen tissue sections¹, immunofluorescent staining², and ELISA³.

Application References:
 1. Ma S, *et al.* 2011. *J. Virol.* 85:165. (IHC)
 2. Manzo A, *et al.* 2008. *Arthritis Rheum.* 11:3377. (IF)
 3. Kato K, *et al.* 2007. *Exp. Hematol.* 35:434. (ELISA)

Description: CD27 is a 50-55 kD type I membrane protein also known as S152 and T14. It is a

lymphocyte-specific member of the TNF-receptor superfamily. CD27 is expressed on medullary thymocytes, virtually all mature T cells, some B cells, and NK cells. CD27 binds to CD70, and plays a role in costimulation of T cell activation and regulation of B cell differentiation and proliferation. The cytoplasmic domains of CD27 have also been shown to interact with TRAF2 and TRAF5 to elicit NF- κ B and SAPK/JNK activation.

**Antigen
References:**

1. Knapp W, *et al.* 1989. Leucocyte Typing IV: White Cell Differentiation Antigens. Oxford University Press.
2. Schlossman S, *et al.* 1995. Leucocyte Typing V: White Cell Differentiation Antigens. Oxford University Press.
- 3.