

Brilliant Violet 421™ anti-human CD4

Catalog # / Size: 2387115 / 25 tests
2387120 / 100 tests

Clone: A161A1

Isotype: Rat IgG2b, κ


Immunogen: Human CD4 T cells

Reactivity: Human

Preparation: The antibody was purified by affinity chromatography and conjugated with Brilliant Violet 421™ under optimal conditions. The solution is free of unconjugated Brilliant Violet 421™ and unconjugated antibody.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and BSA (origin USA).

Concentration: Lot-specific



Human peripheral blood lymphocytes were stained with CD4 (clone A161A1) Brilliant Violet 421™ (filled histogram) or rat IgG2b, κ Brilliant Violet 421™ isotype control (open histogram).

Applications:

Applications: Flow Cytometry

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is 5 µl per million cells or 5 µl per 100 µl of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

Brilliant Violet 421™ excites at 405 nm and emits at 421 nm. The standard bandpass filter 450/50 nm is recommended for detection. Brilliant Violet 421™ is a trademark of Sirigen Group Ltd.

This product is subject to proprietary rights of Sirigen Inc. and is made and sold under license from Sirigen Inc. The purchase of this product conveys to the buyer a non-transferable right to use the purchased product for research purposes only. This product may not be resold or incorporated in any manner into another product for resale. Any use for therapeutics or diagnostics is strictly prohibited. This product is covered by U.S. Patent(s), pending patent applications and foreign equivalents.

- Application References:**
1. Zhu J, *et al.* 2010. *Annu Rev. Immunol.* 28:445.
 2. Vignali DA. 2010. *J. Immunol.* 184:5933.
 3. Zhou L, *et al.* 2009. *Immunity* 30:646.
 4. Singer A, *et al.* 2008. *Nat. Rev. Immunol.* <

Description: CD4, also known as T4/Leu-3, is a 55 kD single-chain type I transmembrane glycoprotein and member of the immunoglobulin superfamily. It is expressed on most thymocytes, helper T cells, type II NKT cells, and monocytes/macrophages. CD4 is part of the TCR/CD3 complex, binds to β2 domain from the MHC class II molecule, and participates in TCR signal transduction. CD4 is the receptor of IL-16

and is a coreceptor for the human immunodeficiency virus (HIV) and human herpes virus 7 (HHV-7).

**Antigen
References:**

1. Zhu J, *et al.* 2010. *Annu Rev. Immunol.* 28:445.
2. Vignali DA. 2010. *J. Immunol.* 184:5933.
3. Zhou L, *et al.* 2009. *Immunity* 30:646.
4. Singer A, *et al.* 2008. *Nat. Rev. Immunol.* <