## SONY

## **Product Data Sheet**

## Brilliant Violet 421™ anti-human ROR1

**Catalog # / Size:** 2389085 / 25 tests

2389090 / 100 tests

Clone: 2A2

**Isotype:** Mouse IgG1, κ

Reactivity: Human

Concentration: Lot-specific

## **Applications:**

**Applications:** Flow Cytometry

Recommended

**Usage:** 

Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is 5 microL per million cells or 5 microL per 100 microL of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

Brilliant Violet  $421^{\text{TM}}$  excites at 405 nm and emits at 421 nm. The standard bandpass filter 450/50 nm is recommended for detection. Brilliant Violet  $421^{\text{TM}}$  is a trademark of Sirigen Group Ltd.

This product is subject to proprietary rights of Sirigen Inc. and is made and sold under license from Sirigen Inc. The purchase of this product conveys to the buyer a non-transferable right to use the purchased product for research purposes only. This product may not be resold or incorporated in any manner into another product for resale. Any use for therapeutics or diagnostics is strictly prohibited. This product is covered by U.S. Patent(s), pending patent applications and foreign equivalents.

Application

Notes:

Clone 2A2 binds to the membrane distal Ig-domain of ROR1. Clone 2A2 has also

been shown to exhibit cross reactivity towards mouse ROR12.

**Description:** 

ROR1, also known as NTRKR1, is a type I transmembrane protein and member of the ROR subfamily of surface receptors. ROR1 consists of one frizzled domain, one Ig-like C2-type domain, one kringle domain, and one kinase domain with no catalytic activity. ROR1 is expressed on embryonic tissue, in the central nervous system and on some cancer cells, and is used as a marker for B-cell chronic lymphocytic leukemia. Wnt5a has been identified as a ligand for ROR1.