Product Data Sheet

PE/Dazzle[™] 594 anti-human ROR1

Catalog # / Size:	2389095 / 25 tests 2389100 / 100 tests	ġ. I
Clone:	2A2	
Isotype:	Mouse lgG1, к	<u>a</u>
Immunogen:	ROR1-Fc fusion protein	
Reactivity:	Human	e e
Preparation:	The antibody was purified by affinity chromatography and conjugated with PE/Dazzle [™] 594 under optimal conditions. The solution is free of unconjugated PE/Dazzle [™] 594 and unconjugated antibody.	0 10 ² 10 ³ 10 ⁴ 10 ⁵ Log Fluorescence Intensity
Formulation:	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA (origin USA).	Human teratocarcinoma cell line NCCIT was stained with ROR1 (clone 2A2) PE/Dazzle™ 594 (filled histogram) or mouse IgG1, ĸ PE/Dazzle™ 594 isotype control (open histogram).
Concentration:	Lot-specific	

Applications:

Applications:	Flow Cytometry	
Recommended Usage:	Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is 5 μ l per million cells or 5 μ l per 100 μ l of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.	
	* PE/Dazzle $^{ m M}$ 594 has a maximum excitation of 566 nm and a maximum emission of 610 nm.	
Application Notes:	Clone 2A2 binds to the membrane distal Ig-domain of ROR1. Clone 2A2 has also been shown to exhibit cross reactivity towards mouse ROR1 ² .	
Application References:	1. Bicocca VT, <i>et al.</i> 2012. <i>Cancer Cell.</i> 22:656.	
	2. Zhang S, <i>et al.</i> 2012. <i>PLoS One</i> 7:e31127.	
	3. Uhrmacher S, <i>et al.</i> 2011. <i>Leuk Res.</i> 35:1360.	
	4. Yang J, <i>et al.</i> 2011.	
Description:	ROR1, also known as NTRKR1, is a type I transmembrane protein and member of the ROR subfamily of surface receptors. ROR1 consists of one frizzled domain, one Ig-like C2-type domain, one kringle domain, and one kinase domain with no catalytic activity. ROR1 is expressed on embryonic tissue, in the central nervous system and on some cancer cells, and is used as a marker for B-cell chronic lymphocytic leukemia. Wnt5a has been identified as a ligand for ROR1.	
Antigen References:	1. Bicocca VT, <i>et al.</i> 2012. <i>Cancer Cell.</i> 22:656.	

For research use only. Not for diagnostic use. Not for resale. Sony Biotechnology Inc. will not be held responsible for patent infringement or other violations that may occur with the use of our products. Sony Biotechnology Inc. 1730 North First Street, San Jose, CA 95112 www.sonybiotechnology.com

- 2. Zhang S, et al. 2012. PLoS One 7:e31127.
- 3. Uhrmacher S, et al. 2011. Leuk Res. 35:1360.
- 4. Yang J, *et al.* 2011.