

PE/Cy7 anti-human CD120b

Catalog # / Size: 2392060 / 100 tests
2392055 / 25 tests

Clone: 3G7A02

Isotype: Rat IgG2a, κ


Immunogen: Partial CD120b recombinant protein expressed in 293E cells (22-211aa).

Reactivity: Human

Preparation: The antibody was purified by affinity chromatography and conjugated with PE/Cy7 under optimal conditions. The solution is free of unconjugated PE/Cy7 and unconjugated antibody.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA (origin USA).

Concentration: Lot-specific



Human peripheral blood lymphocytes were stained with CD120b (clone 3G7A02) PE/Cy7 (filled histogram) or rat IgG2a, κ PE/Cy7 isotype control (open histogram).

Applications:

Applications: Flow Cytometry

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is 5 microL per million cells or 5 microL per 100 microL of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

Description: CD120b, also known as TNFRII or TNFRSF1B, belongs to the TNF receptor superfamily and can interact with ligands TNF-α and TNF-β (also known as LT-α). It is a 75 kD type I transmembrane protein expressed on lymphocytes, monocytes, and granulocytes. When present in a heterocomplex with CD120a, it induces the recruitment of anti-apoptotic molecules, c-IAP1 and c-IAP2 that possess E3 ubiquitin ligase activity. Mutations in CD120b are associated with TNF-associated periodic syndrome (TRAPS) or periodic fever syndrome.

- Antigen References:**
1. Andrews JS, *et al.* 1990. *J. Immunol.* 144:2582.
 2. Nophar Y, *et al.* 1990. *EMBO J.* 9:3269.
 3. Kohno T, *et al.* 1990. *Proc. Natl. Acad. Sci. USA* 87:8331.
 4. Tartaglia LA, <