

PE/Dazzle™ 594 anti-human CD57

Catalog # / Size: 2398095 / 25 tests
2398100 / 100 tests

Clone: HNK-1

Isotype: Mouse IgM, κ


Immunogen: Membrane extract of human lymphoblastoid cell line HSB-2.

Reactivity: Human

Preparation: The antibody was purified by affinity chromatography and conjugated with PE/Dazzle™ 594 under optimal conditions. The solution is free of unconjugated PE/Dazzle™ 594 and unconjugated antibody.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA (origin USA).

Concentration: Lot-specific



Human peripheral blood lymphocytes were stained with CD8 APC and CD57 (clone HNK-1) PE/Dazzle™ 594.

Applications:

Applications: Flow Cytometry

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is 5 microL per million cells or 5 microL per 100 microL of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

* PE/Dazzle™ 594 has a maximum excitation of 566 nm and a maximum emission of 610 nm.

Application Notes: Additional reported applications for the relevant formats include: Western blotting¹.

Application References: 1. Yoshihara Y, *et al.* 1991. *J. Cell Biol.* 115:731. (WB)
2. Abo T, *et al.* 1981. *J. Immunol.* 127:1024.
3. Abo T, *et al.* 1982. *J. Immunol.* 129:1752.
4. Abo T, *et al.* 1982. *J. Immunol.* 129:1758.

Description: CD57, also known as HNK-1, NK-1, and Leu-7 is a 100-115 kD oligosaccharide antigenic determinant expressed on a variety of proteins, lipids, and chondroitin sulfate proteoglycans. CD57 is expressed on a subset of peripheral blood lymphocytes, including NK cells and CD8⁺ T cells, and is also expressed on neural cells and striated muscle. CD57 is not expressed on red blood cells, granulocytes, monocytes, or platelets. While the function of CD57 is unknown, binding to L-selectin, P-selectin, and a fragment of laminin suggests that CD57 may be involved in cell-matrix interactions. CD57 is increased in some disease states associated with CD4/CD8 imbalances (AIDS, autoimmune disease, viral infections, and allograft transplants).

Antigen References: 1. Schubert J, *et al.* 1989. In *Leucocyte Typing IV* (Knapp W, ed) Oxford University Press Oxford pp 711-714.

2. Palmer BE, *et al.* 2005. *J. Immunol.* 175:8415.
3. Schachner M, *et al.* 1995. *Trends Neurosc*