


**PE/Cy7 anti-human CD309 (VEGFR2)**

|                          |  |
|--------------------------|--|
| <b>Catalog # / Size:</b> | 2399555 / 25 tests<br>2399560 / 100 tests  |
| <b>Clone:</b>            | 7D4-6  |
| <b>Isotype:</b>          | Mouse IgG1, $\kappa$   |
| <b>Immunogen:</b>        | Human KDR recombinant protein  |
| <b>Reactivity:</b>       | Human  |
| <b>Preparation:</b>      | The antibody was purified by affinity chromatography and conjugated with PE/Cy7 under optimal conditions. The solution is free of unconjugated PE/Cy7 and unconjugated antibody. |
| <b>Formulation:</b>      | Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA (origin USA).  |
| <b>Concentration:</b>    | Lot-specific   |



HUVEC human endothelial cells were stained with CD309 (clone 7D4-6) PE/Cy7 (filled histogram) or mouse IgG1,  $\kappa$  PE/Cy7 isotype control (open histogram).

**Applications:**

|                           |  |
|---------------------------|--|
| <b>Applications:</b>      | Flow Cytometry   |
| <b>Recommended Usage:</b> | Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is 5 microL per million cells or 5 microL per 100 microL of whole blood. It is recommended that the reagent be titrated for optimal performance for each application. |

This product is subject to proprietary rights of Sirigen Inc. and is made and sold under license from Sirigen Inc. The purchase of this product conveys to the buyer a non-transferable right to use the purchased product for research purposes only. This product may not be resold or incorporated in any manner into another product for resale. Any use for therapeutics or diagnostics is strictly prohibited. This product is covered by U.S. Patent(s), pending patent applications and foreign equivalents.

**Description:** CD309, also known as VEGF-R2, KDR, and Flk-1 (mouse), is a type I transmembrane glycoprotein. It is a member of the CSF-1/PDGF receptor family of type III tyrosine kinase receptors. Human VEGF-R2 is mainly expressed by endothelial cells, embryonic tissues, and megakaryocytes. It plays an important role in the regulation of angiogenesis, vasculogenesis, and vascular permeability. The ligands of VEGF-R2 include VEGF-A, VEGF-C, and VEGF-D splice isoforms. Activation of VEGF-R2 with its ligands results in the receptor dimerization and autophosphorylation, stimulating endothelial cell proliferation and migration.

**Antigen References:**

1. Zola H, *et al.* 2007. Leukocyte and Stromal Cell Molecules: The CD Markers Wiley-Liss A John Wiley & Sons Inc, Publication.
2. Ferrara N and Gerber HP. 2002. *Acta. Haematol.* 106:148.
3. Murohara T, *et al.*