SONY

Product Data Sheet

Brilliant Violet 421™ anti-human CD11a/CD18 (LFA-1)

Catalog # / Size: 2417040 / 100 tests

2417035 / 25 tests

Clone: m24

Isotype: Mouse IgG1, κ

Reactivity: Human

Concentration: Lot-specific

Applications:

Applications: Flow Cytometry

Recommended

Usage:

Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is ≤ 5 microL per million cells or 5 microL per 100 microL of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

Brilliant Violet 421^{TM} excites at 405 nm and emits at 421 nm. The standard bandpass filter 450/50 nm is recommended for detection. Brilliant Violet 421^{TM} is a trademark of Sirigen

This product is subject to proprietary rights of Sirigen Inc. and is made and sold under license from Sirigen Inc. The purchase of this product conveys to the buyer a non-transferable right to use the purchased product for research purposes only. This product may not be resold or incorporated in any manner into another product for resale. Any use for therapeutics or diagnostics is strictly prohibited. This product is covered by U.S. Patent(s), pending patent applications and foreign equivalents.

cqc

Application Notes:

Clone m24 can be used as a reporter of the activation state of integrin receptor in

response to exposure with Ma²⁺ or Mn²⁺.

Description:

CD11/CD18 belongs to the integrin family of proteins. It is heterodimeric cell surface receptor expressed on all leukocytes. CD18, in association with integrin α chain CD11a, CD11b, and CD11c forms LFA-1, Mac-1, and $\alpha_X\beta_2$, respectively, and plays an important role in leukocyte adhesion. CD11/CD18 complexes bind ICAM-1 (CD54), ICAM-2 (CD102), ICAM-3 (CD50), iC3b, and fibrinogen. Clone m24 binds the extended/open high affinity conformation of CD11a/CD18. The antibody can be used as a reporter of the activation state of the integrin receptor in response to exposure to Mg^{2+} or Mn^{2+} .