

APC/Fire™ 750 anti-human CD22

Catalog # / Size: 2417610 / 100 tests
2417605 / 25 tests

Clone: S-HCL-1

Isotype: Mouse IgG2b, κ

Immunogen: Fibronectin-purified human monocytes


Reactivity: Human

Preparation: The antibody was purified by affinity chromatography and conjugated with APC/Fire™ 750 under optimal conditions.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA (origin USA)

Workshop Number: 750 under optimal conditions.

Concentration: Lot-specific




Human peripheral blood lymphocytes were stained with CD19 PE and CD22 (clone S-HCL-1) APC/Fire™ 750 (left) or mouse IgG2b, κ APC/Fire™ 750 isotype control (right).

Applications:

Applications: Flow Cytometry

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is 5 µL per million cells in 100 µL staining volume or 5 µL per 100 µL of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.



* APC/Fire™ 750 has a maximum excitation of 650 nm and a maximum emission of 787 nm.

- Application Notes:**
1. Nitschke L. 2005. *Curr. Opin. Immunol.* 17:290
 2. Foon Ka, et al. 1986. *Blood.* 68:297
 3. Schwarting R, et al. 1985. *Blood.* 65:974
 4. Campana D, et al. 1985. *J. Immunol.* 134:1524

- Application References:**
1. Hogg N. et al. 1985. *Cell Immunol.* 92:247.
 2. McDowall A. et al. 2003. *J. Clin. Invest.* 111:51.

Description: CD22 is a 130 kD type I transmembrane glycoprotein also known as Siglec-2 and BL-CAM and is a member of the immunoglobulin superfamily (sialoadhesion subgroup). CD22 is expressed in the cytoplasm of pro-B and pre-B cells, and on the surface of mature B and activated B cells, but not on plasma cells. CD22 is present in the B cell receptor complex and associates with SHP-1, Syk, Lck, Lyn, and phospholipase C γ 1. A primary function of CD22 is thought to be in limiting antigen receptor signaling by modulating B cell activation threshold. CD22 has been shown to bind to CD45RO and CD75, although the natural ligands for this molecule remain controversial.

Antigen 1. Clark E. 1993. *J. Immunol.* 150:4715.
References: 2. Shan D, *et al.* 1995. *J. Immunol.* 154:4466.