

Brilliant Violet 421™ anti-human CD5

Catalog # / Size: 2420145 / 25 tests
2420150 / 100 tests

Clone: L17F12

Isotype: Mouse IgG2a, κ

Immunogen: Human T-acute lymphoblastic leukemia (ALL) cells


Reactivity: Human

Preparation: The antibody was purified by affinity chromatography and conjugated with Brilliant Violet 421™ under optimal conditions. The solution is free of unconjugated Brilliant Violet 421™ and unconjugated antibody.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and BSA (origin USA).

Workshop Number: II T7

Concentration: Lot-specific



Human peripheral blood lymphocytes were stained with Brilliant Violet 421™ anti-human CD5 (clone L17F12, filled histogram) or Brilliant Violet 421™ mouse IgG2a, κ isotype control (open histogram).

Applications:

Applications: Flow Cytometry

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is 5 µl per million cells or 5 µl per 100 µl of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

Brilliant Violet 421™ excites at 405 nm and emits at 421 nm. The standard bandpass filter 450/50 nm is recommended for detection. Brilliant Violet 421™ is a trademark of Sirigen Group Ltd.

This product is subject to proprietary rights of Sirigen Inc. and is made and sold under license from Sirigen Inc. The purchase of this product conveys to the buyer a non-transferable right to use the purchased product for research purposes only. This product may not be resold or incorporated in any manner into another product for resale. Any use for therapeutics or diagnostics is strictly prohibited. This product is covered by U.S. Patent(s), pending patent applications and foreign equivalents.

Application Notes: Additional reported applications (for the relevant formats): immunoprecipitation^{1,3,4} and immunohistochemical staining of frozen and formalin-fixed paraffin-embedded sections^{2,3,5}.

- Application References:**
1. Kipps T. 1989. *Adv. Immunol.* 47:117.
 2. Resnick D, et al. 1994. *Trends Biochem. Sci.* 1:5.
 3. Wood GS and Freudenthal PS. 1992. *Am. J. Pathol.* 141:789.

Description: CD5, also known as Leu-1, Ly-1 and T1, is a 67 kD single chain type I glycoprotein that is a member of the scavenger receptor superfamily. CD5 is expressed on T cells, thymocytes, B cell subsets, chronic B lymphocytic leukemia (B-Cells), and peripheral blood dendritic cells. CD5 modulates T and B cell receptor signaling, thymocyte maturation, and T-B cell interactions. One of its ligands is CD72.

**Antigen
References:**

1. Kipps T. 1989. *Adv. Immunol.* 47:117.
2. Resnick D, *et al.* 1994. *Trends Biochem. Sci.* 1:5.
3. Wood GS and Freudenthal PS. 1992. *Am. J. Pathol.* 141:789.