

APC/Cy7 anti-human CD33

Catalog # / Size: 2433070 / 100 tests
2433065 / 25 tests

Clone: P67.6

Isotype: Mouse IgG1, κ


Immunogen: FMY9S5 cells expressing CD33 gene.

Reactivity: Human

Preparation: The antibody was purified by affinity chromatography and conjugated with APC/Cy7 under optimal conditions. The solution is free of unconjugated APC/Cy7 and unconjugated antibody.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA (origin USA).

Concentration: Lot-specific



Human peripheral blood monocytes were stained with CD33 (clone P67.6) APC/Cy7 (filled histogram) or mouse IgG1, κ APC/Cy7 isotype control (open histogram).

Applications:

Applications: Flow Cytometry

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is 5 microL per million cells or 5 microL per 100 microL of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

Application References: 1. Hoyer J, *et al.* 2008. *Am. J. Clin. Pathol.* 129:316.

Description: CD33, also known as Siglec-3, gp67, and p67, is a 67 kD type I transmembrane glycoprotein. It is a sialoadhesion immunoglobulin superfamily member, which is expressed on myeloid progenitors, monocytes, granulocytes, dendritic cells, and mast cells. CD33 is absent on normal platelets, lymphocytes, erythrocytes, and hematopoietic stem cells. CD33 functions as a sialic acid-dependent cell adhesion molecule with carbohydrate/lectin binding activity.

Antigen References: 1. Favaloro E, *et al.* 1988. *Br. J. Haematol.* 69:163.
2. Freeman S, *et al.* 1995. *Blood* 85:2005.