

PE/Fire™ 700 anti-human CD14

Catalog # / Size: 2435785 / 25 tests
2435790 / 100 tests

Clone: 63D3

Isotype: Mouse IgG1, κ


Immunogen: Purified human peripheral blood monocytes.

Reactivity: Human

Preparation: The antibody was purified by affinity chromatography and conjugated with PE/Fire™ 700 under optimal conditions.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA (origin USA)

Concentration: Lot-specific



Human peripheral blood monocytes were stained with anti-human HLA-DR FITC and anti-human CD14 PE/Fire™ 700 (clone 63D3) (left), or mouse IgG1, κ PE/Fire™ 700 (right).

Applications:

Applications: Flow Cytometry

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is 5 µL per million cells in 100 µL staining volume or 5 µL per 100 µL of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

* PE/Fire™ 700 has a maximum excitation of 565 nm and a maximum emission of 695 nm.

Application References: 1. Fridlender ZG, *et al.* 1999. *Hum. Immunol.* 11:1028.
2. Devitt A, *et al.* 1998. *Nature* 6675:505.

Description: CD14 is a 53-55 kD glycosylphosphatidylinositol (GPI)-linked membrane glycoprotein that is also known as the LPS receptor. CD14 is expressed at high levels on monocytes and macrophages, and at lower levels on granulocytes. Some dendritic cell populations such as interfollicular dendritic cells, reticular dendritic cells, and Langerhans cells have also been reported to express CD14. As a high-affinity receptor for LPS, CD14 is involved in the clearance of gram-negative pathogens and in the upregulation of adhesion molecules and cytokine expression in monocytes and neutrophils.

Antigen References: 1. Stocks SC, *et al.* 1990. *Biochem. J.* 268:275.
2. Wright SD, *et al.* 1990. *Science* 4975:1431.