Product Data Sheet

Purified anti-human CD45

Catalog # / Size: 2442510 / 100 μg

Clone: 2D1

Isotype: Mouse IgG1, κ **Immunogen:** Human PBMC

Reactivity: Human


Preparation: The antibody was purified by affinity

chromatography.

Formulation: Phosphate-buffered solution, pH 7.2,

containing 0.09% sodium azide.

Concentration: Lot-specific



Human peripheral blood lymphocytes were stained with purified CD45 (clone 2D1) (filled histogram) or purified mouse IgG1, κ isotype control (open histogram), followed by anti-mouse IgG FITC.

Applications:

Applications: Immunofluorescence

Recommended

Usage:

Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of

this reagent is ≤0.5 microg per million cells in 100 microL volume. It is recommended that the reagent be titrated for optimal performance for each

application.

Application Notes:

It was found that the HI30 clone and the 2D1 clone can cross block each other's

binding.

Application

1. Bradstock KF, et al. 1980. J. Natl. Cancer Inst. 65:33.

References: 2. Csib

2. Csiba A, et al. 1984. Br. J. Cancer 50:699.

3. Tchilian EZ, et al. 2001. J. Immunol. 166:1308.

4. Lee MS, et al. 2004. Int. Immunol. 16:1109.

Description:

CD45 is a 180 - 240 kD single chain type I membrane glycoprotein also known as leukocyte common antigen (LCA) and T200. It is a tyrosine phosphatase

expressed on the plasma membrane of all hematopoietic cells,

except erythrocytes or platelets. CD45 is a signaling molecule that regulates a variety of cellular processes including cell growth, differentiation, cell cycle, and oncogenic transformation. CD45 plays a critical role in T and B cell antigen receptor-mediated activation by dephosphorylating substrates including p56Lck, p59Fyn, and other Src family kinases. CD45 non-covalently associates with lymphocyte phosphatase-associated phosphoprotein (LPAP) on T and B lymphocytes. CD45 has been reported to bind galectin-1 and to be associated with several other cell surface antigens including CD1, CD2, CD3, and CD4.

Antigen References: 1. Thomas M. 1989. Annu. Rev. Immunol. 7:339.

References: 2. Trowbridge I, et al. 1994. Annu. Rev. Immunol. 12:85.