

FITC anti-human CD45

Catalog # / Size: 2442535 / 25 tests
2442540 / 100 tests

Clone: 2D1

Isotype: Mouse IgG1, κ


Immunogen: Human PBMC

Reactivity: Human

Preparation: The antibody was purified by affinity chromatography and conjugated with FITC under optimal conditions. The solution is free of unconjugated FITC and unconjugated antibody.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA (origin USA).

Concentration: Lot-specific



Human peripheral blood lymphocytes were stained with CD45 (clone 2D1) FITC (filled histogram) or mouse IgG1, κ FITC isotype control (open histogram).

Applications:

Applications: Flow Cytometry

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is 5 microL per million cells or 5 microL per 100 microL of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

Application Notes: It was found that the HI30 clone and the 2D1 clone can cross block each other's binding.

- Application References:**
1. Bradstock KF, *et al.* 1980. *J. Natl. Cancer Inst.* 65:33.
 2. Csiba A, *et al.* 1984. *Br. J. Cancer* 50:699.
 3. Tchilian EZ, *et al.* 2001. *J. Immunol.* 166:1308.
 4. Lee MS, *et al.* 2004. *Int. Immunol.* 16:1109.

Description: CD45 is a 180 - 240 kD single chain type I membrane glycoprotein also known as leukocyte common antigen (LCA) and T200. It is a tyrosine phosphatase expressed on the plasma membrane of all hematopoietic cells, except erythrocytes or platelets. CD45 is a signaling molecule that regulates a variety of cellular processes including cell growth, differentiation, cell cycle, and oncogenic transformation. CD45 plays a critical role in T and B cell antigen receptor-mediated activation by dephosphorylating substrates including p56Lck, p59Fyn, and other Src family kinases. CD45 non-covalently associates with lymphocyte phosphatase-associated phosphoprotein (LPAP) on T and B lymphocytes. CD45 has been reported to bind galectin-1 and to be associated with several other cell surface antigens including CD1, CD2, CD3, and CD4.

- Antigen References:**
1. Thomas M. 1989. *Annu. Rev. Immunol.* 7:339.
 2. Trowbridge I, *et al.* 1994. *Annu. Rev. Immunol.* 12:85.