
Product Data Sheet

Alexa Fluor® 647 anti-human CD223 (LAG-3)


Catalog # / Size:	2446015 / 25 tests 2446020 / 100 tests
Clone:	7H2C65
Isotype:	Mouse IgG1, к
Immunogen:	Human LAG-3 transfected cells.
Reactivity:	Human
Preparation:	The antibody was purified by affinity chromatography and conjugated with Alexa Fluor® 647 under optimal conditions.
Formulation:	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA (origin USA).
Concentration:	0.5



CD3/CD28/IL-2 stimulated (three days) peripheral blood monocular cells (PBMCs) were stained with CD8 FITC and CD223 (clone 7H2C65) Alexa Fluor® 647 (top) or mouse IgG1, κ Alexa Fluor® 647 isotype control (bottom).

Applications:

		0
Applications:	Flow Cytometry	Contr
Recommended Usage:	Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is 5 microL per million cells or 5 microL per 100 microL of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.	Mouse IgG1, x Isotype Control 6 6 6 6
	* Alexa Fluor® 647 has a maximum emission of 668 nm when it is excited at 633 nm / 635 nm.	
Application Notes:	The staining of clone 7H2C65 cannot be blocked by clone 11C3C65, which is another anti-human CD223 (LAG-3) antibody.	
Description:	CD223, also known as LAG-3, is a 70 kD ty is involved in T-cell signaling. Similar to CE a higher affinity. CD223 negatively regulat activated T-cells and natural killer cells (N	04, CD tes T-o



Description: CD223, also known as LAG-3, is a 70 kD type I transmembrane glycoprotein that is involved in T-cell signaling. Similar to CD4, CD223 binds MHC class II, but with a higher affinity. CD223 negatively regulates T-cell activation. It is expressed by activated T-cells and natural killer cells (NKs), as well as regulatory T-cells. It is transiently expressed on the surface of activated T-cells in acute conditions but high expression is maintained under tolerizing conditions. CD223 deficiency results in reduced tumor growth. CD223 and PD-1 can act in synergy and reverse exhausted phenotypes, improve tumor rejection, and control viral load.

For research use only. Not for diagnostic use. Not for resale. Sony Biotechnology Inc. will not be held responsible for patent infringement or other violations that may occur with the use of our products. Sony Biotechnology Inc. 1730 North First Street, San Jose, CA 95112 www.sonybiotechnology.com

 Castelli C, *et al.* 2014. *Oncoimmunology* 3(11):e967146.
Poirier N, *et al.* 2011. *Clin. Exp. Immunol.* 164:265.
Juno JA, *et al.* 2015. *Retrovirology* 12:17.
Casati C, *et al.* Antigen **References:**