Product Data Sheet

APC anti-human CD99

Catalog # / Size:	2456540 / 100 tests 2456535 / 25 tests	
Clone:	3B2/TA8	Belative Cell Number
Isotype:	Mouse lgG1, κ	
Immunogen:	Human thymus	
Reactivity:	Human	
Preparation:	The antibody was purified by affinity chromatography and conjugated with APC under optimal conditions. The solution is free of unconjugated APC and unconjugated antibody.	
Formulation:	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA (origin USA).	Log Fluorescence Intensity Human peripheral blood lymphocytes were stained with CD99 APC (clone 3B2/TA8, filled histogram) or mouse IgG1, κ APC isotype control (open histogram).
Concentration:	Lot-specific	

Applications:

Applications:Flow CytometryRecommended
Usage:Each lot of this antibody is quality control tested by immunofluorescent staining
with flow cytometric analysis. For flow cytometric staining, the suggested use of
this reagent is 5 microL per million cells or 5 microL per 100 microL of whole
blood. It is recommended that the reagent be titrated for optimal performance for
each application.

 Application
 1. Waclavicek M, et al. 1998. J. Immunol. 161:4671.

 References:
 2. Pickl W, et al. 2001. J. Virol. 75:7175.

Description: CD99 is a type I single chain transmembrane protein devoid of N-linked glycosylation sites encoded by the pseudoautosomal gene MIC2. CD99 has an apparent molecular weight of 32 kD and is widely expressed on a variety of tissues. CD99 is highly expressed on thymocytes, T cells, and T cell leukemias and lymphomas. However, it is absent on some B cell lines, fetal B cells, eosinophils, granulocytes and the NK-cell line YT. CD99 is involved in spontaneous rosette formation with erythrocytes and may also be involved in other T-cell and hematopoietic cell adhesion pathways. CD99 has been reported to activate a caspase-independent death pathway in T cells under some conditions. CD99 interacts with a number of proteins including ferritin heavy chain 1, karyopherin β 1, TRIP13, cyclophilin A, annexin II, and ubiquitinconjugating enzyme E2H.

Antigen	1. Gelin C, <i>et al.</i> 1989. <i>EMBO.</i> 8:3253.	
References:	2. Goodfellow PJ, <i>et al.</i> 1986. <i>Science</i> 234:740.	
	3. Pettersen RD, et al. 2001. J. Immunol. 166:4931.	

For research use only. Not for diagnostic use. Not for resale. Sony Biotechnology Inc. will not be held responsible for patent infringement or other violations that may occur with the use of our products. Sony Biotechnology Inc. 1730 North First Street, San Jose, CA 95112 www.sonybiotechnology.com