## **Product Data Sheet**

## **APC anti-human CD11c**

Catalog # / Size:	2457525 / 25 tests 2457530 / 100 tests	ά <b>Λ</b>
Clone:	S-HCL-3	
Isotype:	Mouse lgG2b, κ	
Immunogen:	Spleen cells from patient diagnosed with hairy cell leukemia.	
<b>Reactivity:</b>	Human	el attive
Preparation:	The antibody was purified by affinity chromatography and conjugated with APC under optimal conditions. The solution is free of unconjugated APC and unconjugated antibody.	0 10 <sup>3</sup> 10 <sup>4</sup> 10 <sup>5</sup> Log Fluorescence Intensity
Formulation:	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA (origin USA).	Human peripheral blood granulocytes were stained with CD11c (clone S-HCL-3) APC (filled
Concentration:	Lot-specific	histogram) or mouse lgG2b, κ APC isotype control (open histogram).

## **Applications:**

Applications: Recommended Usage: Application	Flow Cytometry Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is 5 microL per million cells or 5 microL per 100 microL of whole blood. It is recommended that the reagent be titrated for optimal performance for each application. Additional reported applications (for the relevant formats) include:
Notes: Application References:	<ol> <li>immunohistochemistry on frozen tissue sections<sup>1,2,3,4</sup> and immunoprecipitation1.</li> <li>Schwarting R, <i>et al.</i> 1985. <i>Blood</i> 65:974.</li> <li>Knowles DM, <i>et al.</i> 1990. <i>Am. J. Pathol.</i> 136:29.</li> <li>Vandenabeele S, <i>et al.</i> 2001. <i>Blood</i> 97:1733.</li> <li>Shaw JL, <i>et al.</i> 2011. <i>J. Reprod. Immunol.</i> 89:84.</li> </ol>
Description: Antigen	CD11c is a 145-150 kD type I transmembrane glycoprotein also known as integrin $\alpha_x$ and CR4. CD11c non-covalently associates with integrin $\beta_2$ (CD18) and is expressed on monocytes/macrophages, dendritic cells, granulocytes, NK cells, and subsets of T and B cells. CD11c has been reported to play a role in adhesion and CTL killing through its interactions with fibrinogen, CD54, and iC3b. 1. Petty HR, Todd RF 3rd. 1996. <i>Immunol. Today</i> 17:209.
References:	2. Springer T. 1994. <i>Cell</i> 76:301. 3. Ihanus E, <i>et al.</i> 2007. <i>Blood</i> 109:802-10.

For research use only. Not for diagnostic use. Not for resale. Sony Biotechnology Inc. will not be held responsible for patent infringement or other violations that may occur with the use of our products. Sony Biotechnology Inc. 1730 North First Street, San Jose, CA 95112 www.sonybiotechnology.com