
Product Data Sheet

PE/Cy7 anti-human CD11c

Catalog # / Size:	2457540 / 100 tests 2457535 / 25 tests		6
Clone:	S-HCL-3		11
Isotype:	Mouse lgG2b, κ	Relative Cell Number	11
Immunogen:	Spleen cells from patient diagnosed with hairy cell leukemia.		
Reactivity:	Human		1 1
Preparation:	The antibody was purified by affinity chromatography and conjugated with PE/Cy7 under optimal conditions. The solution is free of unconjugated PE/Cy7 and unconjugated antibody.		10 ³ Log Fb
Formulation:	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA (origin USA).	Human periph granulocytes CD11c (clone	
Concentration:	Lot-specific	histogram) or PE/Cy7 isotyp	



Human peripheral blood granulocytes were stained with CD11c (clone S-HCL-3) PE/Cy7 (filled histogram) or mouse IgG2b, κ PE/Cy7 isotype control (open histogram).

Applications:

Applications:	Flow Cytometry	
Recommended Usage:	Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is 5 microL per million cells or 5 microL per 100 microL of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.	
Application Notes:	Additional reported applications (for the relevant formats) include: immunohistochemistry on frozen tissue sections ^{1,2,3,4} and immunoprecipitation1.	
Application References:	 Schwarting R, <i>et al.</i> 1985. <i>Blood</i> 65:974. Knowles DM, <i>et al.</i> 1990. <i>Am. J. Pathol.</i> 136:29. Vandenabeele S, <i>et al.</i> 2001. <i>Blood</i> 97:1733. Shaw JL, <i>et al.</i> 2011. <i>J. Reprod. Immunol.</i> 89:84. 	
Description:	CD11c is a 145-150 kD type I transmembrane glycoprotein also known as integrin α_x and CR4. CD11c non-covalently associates with integrin β_2 (CD18) and is expressed on monocytes/macrophages, dendritic cells, granulocytes, NK cells, and subsets of T and B cells. CD11c has been reported to play a role in adhesion and CTL killing through its interactions with fibrinogen, CD54, and iC3b.	
Antigen References:	1. Petty HR, Todd RF 3rd. 1996. <i>Immunol. Today</i> 17:209. 2. Springer T. 1994. <i>Cell</i> 76:301. 3. Ihanus E, <i>et al.</i> 2007. <i>Blood</i> 109:802-10.	

For research use only. Not for diagnostic use. Not for resale. Sony Biotechnology Inc. will not be held responsible for patent infringement or other violations that may occur with the use of our products. Sony Biotechnology Inc. 1730 North First Street, San Jose, CA 95112 www.sonybiotechnology.com